



# Entomological Notes

Department of Entomology

## BOXWOOD PSYLLID

*Cacopsylla busi* (Linnaeus)

The boxwood psyllid is a common pest of boxwood, *Buxus* spp. It is not considered as destructive as other boxwood pests. American boxwood *B. sempervirens* appear to be most susceptible to this pest.

### DESCRIPTION

The eggs are small, orange, and spindle-shaped. Nymphs are covered with a white waxy secretion, which readily distinguishes them from other insects that attack boxwood. Adults are light green insects that are about 3 mm long. Both nymphs and adults have piercing-sucking mouthparts.

### LIFE HISTORY

This species overwinters as eggs. They are laid between bud scales of the host plant during early summer. Eggs start hatching as soon as buds begin to open in early spring. Young nymphs immediately begin feeding by removing plant fluids from tender foliage. Small nymphs develop on expanding foliage. Leaves become cupped and several nymphs may be enclosed in a pocket of foliage. Nymphs usually mature into adults by early June. After mating, females deposit eggs that overwinter on the host plant. One generation occurs each year in Pennsylvania.

### DAMAGE

Feeding damage is very noticeable due to leaf cupping that young nymphs produce on host plants. The leaf cupping results from injury done to leaf tissue as it is developing in rapidly growing leaves. Occasionally, young twig growth is affected by this species.

### MANAGEMENT

Treat affected host plants with registered insecticides when nymphs are present in early May.

## WARNING

Pesticides are poisonous. Read and follow directions and safety precautions on labels. Handle carefully and store in original labeled containers out of the reach of children, pets, and livestock. Dispose of empty containers right away, in a safe manner and place. Do not contaminate forage, streams, or ponds.

Gregory A. Hoover  
Sr. Extension Associate  
Dept. of Entomology  
November 2001

TS-8B

© The Pennsylvania State University 2001

This publication is available in alternative media on request.

Where trade names are used, no discrimination is intended and no endorsement by The Pennsylvania State University or Pennsylvania Department of Agriculture is implied.

Entomological Notes are intended to serve as a quick reference guide and should not be used as a substitute for product label information. Although every attempt is made to produce Entomological Notes that are complete, timely, and accurate, the pesticide user bears the responsibility of consulting the pesticide label and adhering to those directions.

Issued in furtherance of Cooperative Extension Works, Acts of Congress May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture and the Pennsylvania Legislature. T.R. Alter, Director of Cooperative Extension, The Pennsylvania State University.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. It is the policy of the University to maintain an academic and work environment free of discrimination, including harassment. The Pennsylvania State University prohibits discrimination and harassment against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, or veteran status. Discrimination or harassment against faculty, staff, or students will not be tolerated at The Pennsylvania State University. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 201 Willard Building, University Park, PA 16802-2801, Tel 814-865-4700/V, 814-863-1150/TTY.

