



NATIVE HOLLY LEAFMINER

Phytomyza ilicicola Loew

Ornamental plantings of holly, *Ilex* spp., in Pennsylvania are often damaged by leafmining pests. One species is the native holly leafminer that only feeds on the foliage of American holly, *Ilex opaca*, and its cultivars. Other leafminers of plants in the genus *Ilex* include the holly leafminer, *P. ilicis* Curtis, a European species that feeds primarily on English holly, *I. aquifolium*. Additionally, the “inkberry leafminer”, *P. glabricola*, infests inkberry, *I. glabra*. The information contained in this fact sheet pertains to the native holly leafminer.

DESCRIPTION

Eggs hatch into tiny maggots or larvae within the leaf mine. The larvae are yellow and when fully grown are 1.5 mm long. The larval stage of the European species, holly leafminer, resembles the native holly leafminer. Adults are small, 0.8 mm-1.6 mm long, black flies.

LIFE HISTORY

This species overwinters as larvae within the leaf mine. In some cases 12 or more larvae have been found in an individual leaf, but the average number found in a leaf is three. During the spring these maggots pupate. Adults emerge from leaves in May, mate, and females deposit eggs in small slits on the underside of newly developing leaves. The eggs hatch into larvae that start to feed, creating new leaf mines. There is only one generation produced each year.

DAMAGE

American holly foliage is damaged by the feeding activity of larvae that destroy leaf tissue between the upper and lower leaf surfaces. Damage first shows up as slender leaf mines that are seldom over 1.3 cm in length. With continuous feeding, leaf mines broaden and leaves become marred with yellow serpentine or blotch mines (Fig. 1). Heavily infested plants may prematurely drop their leaves and be almost devoid of leaves until new growth begins the following spring.

Females may damage foliage by inserting their ovipositors into leaves. The female jabs the leaf with her ovipositor and plant fluid starts to flow from this wound.




Figure 1. Serpentine blotch mines in a holly leaf.

Both females and males feed on the plant fluid that exudes from the wound. Leaves seriously damaged by these ovipositor punctures often become stunted and badly distorted. Be careful not to confuse this type of damage with leaves mechanically injured by overlapping holly leaf spines.

MANAGEMENT

Infested holly leaves may drop to the ground before adults emerge. Leaves that have dropped prematurely should be gathered and destroyed. This helps reduce adult populations that may attack host plants in the spring.

Registered insecticides should be applied according to label directions to the foliage in early May.

Registered systemic insecticide formulations may also be effective against this pest when applied according to label directions as soil injections early in the growing season. When using these insecticides be sure adequate soil moisture exists so that optimal uptake of these products is achieved.

WARNING

Pesticides are poisonous. Read and follow directions and safety precautions on labels. Handle carefully and store in original labeled containers out of the reach of

children, pets, and livestock. Dispose of empty containers right away, in a safe manner and place. Do not contaminate forage, streams, or ponds.

Gregory A. Hoover
Sr. Extension Associate
Dept. of Entomology
April 2003

TS-21

©The Pennsylvania State University 2003

This publication is available in alternative media on request.

Where tradenames are used, no discrimination is intended and no endorsement by The Pennsylvania State University or Pennsylvania Department of Agriculture is implied.

Entomological Notes are intended to serve as a quick reference guide and should not be used as a substitute for product label information. Although every attempt is made to produce Entomological Notes that are complete, timely, and accurate, the pesticide user bears the responsibility of consulting the pesticide label and adhering to those directions.

Issued in furtherance of Cooperative Extension Works, Acts of Congress May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture and the Pennsylvania Legislature. T.R. Alter, Director of Cooperative Extension, The Pennsylvania State University.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. It is the policy of the University to maintain an academic and work environment free of discrimination, including harassment. The Pennsylvania State University prohibits discrimination and harassment against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, or veteran status. Discrimination or harassment against faculty, staff, or students will not be tolerated at The Pennsylvania State University. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 328 Bouke Building, University Park, PA 16802-5901, Tel 814-865-4700/V, 814-863-1150/TTY.