

```

Clear[ $\beta$ ,  $\gamma$ ,  $\lambda$ ]
(*The model is -- eqn 12-14 in Bjornstad et al.
  N0 (t) = Exp ( $\zeta$ ) S (t) => N0 (t) = Exp ( $\zeta$ ) ( $\lambda$  N1 (t-1) +  $\lambda$  N2 (t-1) +  $\lambda$  N3 (t-1))
  N1 (t) = N0 (t-1) Exp[-  $\beta$  Log[N0 (t-1)] -  $\beta$  Log[N1 (t-1)]]
  N2 (t) =  $\lambda$  N1 (t-1)
  N3 (t) =  $\lambda$  N2 (t-1)
  N4 (t) =  $\lambda$  N3 (t-1)
  S (t) = N2 (t)+N3 (t)+N4 (t) =  $\lambda$  N1 (t-1) +  $\lambda$  N2 (t-1) +  $\lambda$  N3 (t-1)
*)

(*Let x0 = Log[N0], x1 = Log[N1], etc
  The following is then the model re-written in log-abundances: *)
f0 = Exp[ $\zeta$ ] ( $\lambda$  Exp[x1] +  $\lambda$  Exp[ x2] +  $\lambda$  Exp[x3]);
f1 = Exp[x0] Exp[- $\beta$  x0 -  $\gamma$  x1];
f2 =  $\lambda$  Exp[x1];
f3 =  $\lambda$  Exp[x2];
(*f4= $\lambda$  Exp[x3] is not needed*)
fs =  $\lambda$  Exp[x1] +  $\lambda$  Exp[ x2] +  $\lambda$  Exp[x3];

(* Differentiating the log-model with respect to log-abundances (see eqn A4): *)
j11 = D[Log[f0], x0]; j12 = D[Log[f0], x1];
j13 = D[Log[f0], x2]; j14 = D[Log[f0], x3]; j15 = D[Log[f0], s];
j21 = D[Log[f1], x0]; j22 = D[Log[f1], x1];
j23 = D[Log[f1], x2]; j24 = D[Log[f1], x3]; j25 = D[Log[f1], s];
j31 = D[Log[f2], x0]; j32 = D[Log[f2], x1];
j33 = D[Log[f2], x2]; j34 = D[Log[f2], x3]; j35 = D[Log[f2], s];
j41 = D[Log[f3], x0]; j42 = D[Log[f3], x1];
j43 = D[Log[f3], x2]; j44 = D[Log[f3], x3]; j45 = D[Log[f3], s];
j51 = D[Log[fs], x0]; j52 = D[Log[fs], x1];
j53 = D[Log[fs], x2]; j54 = D[Log[fs], x3]; j55 = D[Log[fs], s];

(* Defining the Jacobian for the log-linearized system *)
J = {{j11, j12, j13, j14, j15}, {j21, j22, j23, j24, j25},
     {j31, j32, j33, j34, j35}, {j41, j42, j43, j44, j45}, {j51, j52, j53, j54, j55}};
MatrixForm[J]

```

$$\begin{pmatrix}
 0 & \frac{e^{x_1} \lambda}{e^{x_1} \lambda + e^{x_2} \lambda + e^{x_3} \lambda} & \frac{e^{x_2} \lambda}{e^{x_1} \lambda + e^{x_2} \lambda + e^{x_3} \lambda} & \frac{e^{x_3} \lambda}{e^{x_1} \lambda + e^{x_2} \lambda + e^{x_3} \lambda} & 0 \\
 1 - \beta & -\gamma & 0 & 0 & 0 \\
 0 & 1 & 0 & 0 & 0 \\
 0 & 0 & 1 & 0 & 0 \\
 0 & \frac{e^{x_1} \lambda}{e^{x_1} \lambda + e^{x_2} \lambda + e^{x_3} \lambda} & \frac{e^{x_2} \lambda}{e^{x_1} \lambda + e^{x_2} \lambda + e^{x_3} \lambda} & \frac{e^{x_3} \lambda}{e^{x_1} \lambda + e^{x_2} \lambda + e^{x_3} \lambda} & 0
 \end{pmatrix}$$

(* simplifying at equilibrium *)

```
Jeq = Simplify[J /. Exp[x3] -> λ Exp[x2] /. Exp[x2] -> λ Exp[x1]];
MatrixForm[Jeq]
```

$$\begin{pmatrix} 0 & \frac{1}{1+\lambda+\lambda^2} & \frac{\lambda}{1+\lambda+\lambda^2} & \frac{\lambda^2}{1+\lambda+\lambda^2} & 0 \\ 1-\beta & -\gamma & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & \frac{1}{1+\lambda+\lambda^2} & \frac{\lambda}{1+\lambda+\lambda^2} & \frac{\lambda^2}{1+\lambda+\lambda^2} & 0 \end{pmatrix}$$

(* Defining the A vector (cf eqn A5) *)

```
a1 = D[Log[f0], ζ]; a2 = D[Log[f1], ζ];
a3 = D[Log[f3], ζ]; a4 = D[Log[f4], ζ]; a5 = D[Log[fs], ζ];
A = Transpose[{a1, a2, a3, a4, a5}];
MatrixForm[A]
```

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

(*Calculating the vector of transfer functions -- eqn 5 in Bjornstad et al.

NOTE: the below is cast in terms of angular frequencies, w: w = 2 π f*)

```
Id := IdentityMatrix[5];
F := Inverse [Id - Exp[-I w] Jeq] . A
MatrixForm[F]
```

$$\begin{pmatrix} \frac{1+e^{-i w} \gamma}{1+e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \\ \frac{e^{-i w} - e^{-i w} \beta}{1+e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \\ \frac{e^{-2 i w} - e^{-2 i w} \beta}{1+e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \\ \frac{e^{-3 i w} - e^{-3 i w} \beta}{1+e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \\ - \frac{e^{-i w} (1-\beta) \left(\frac{e^{-i w}}{1+\lambda+\lambda^2} - \frac{e^{-2 i w} \lambda}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda^2}{1+\lambda+\lambda^2} \right)}{1+e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \end{pmatrix}$$

```
T := ExpToTrig[F]
PS := Re[T] ^ 2 + Im[T] ^ 2;
```

(*The transfer function for age 0 cohort*)

```
F[[1]]
```

$$\left\{ \frac{1+e^{-i w} \gamma}{1+e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \right\}$$

(*delay coordinate AR-coefficients for young-of-the-year*)
 Simplify[Coefficient[-Denominator[F[[1]]][[1]], {e^{-i w}, e^{-2 i w}, e^{-3 i w}, e^{-4 i w}}]
 (*These are the x0 (t-1), ..., x0 (t-4) coefficients*)

$$\left\{ -\gamma, \frac{1-\beta}{1+\lambda+\lambda^2}, \frac{\lambda-\beta\lambda}{1+\lambda+\lambda^2}, -\frac{(-1+\beta)\lambda^2}{1+\lambda+\lambda^2} \right\}$$

(*AND delay coordinate MA-coefficients for young-of-the-year*)
 Simplify[Coefficient[Numerator[F[[1]]][[1]], {e^{-i w}, e^{-2 i w}, e^{-3 i w}, e^{-4 i w}}]
 (*These are the α (t), ..., α (t-3) coefficients*)

$$\{\gamma, 0, 0, 0\}$$

(*The transfer function for age 1 cohort*)
 F[[2]]

$$\left\{ \frac{e^{-i w} - e^{-i w} \beta}{1 + e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \right\}$$

(*The transfer function for spawning stock cohort*)
 F[[5]]

$$\left\{ -\frac{e^{-i w} (1-\beta) \left(-\frac{e^{-i w}}{1+\lambda+\lambda^2} - \frac{e^{-2 i w} \lambda}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda^2}{1+\lambda+\lambda^2} \right)}{1 + e^{-i w} \gamma - \frac{e^{-2 i w}}{1+\lambda+\lambda^2} + \frac{e^{-2 i w} \beta}{1+\lambda+\lambda^2} - \frac{e^{-3 i w} \lambda}{1+\lambda+\lambda^2} + \frac{e^{-3 i w} \beta \lambda}{1+\lambda+\lambda^2} - \frac{e^{-4 i w} \lambda^2}{1+\lambda+\lambda^2} + \frac{e^{-4 i w} \beta \lambda^2}{1+\lambda+\lambda^2}} \right\}$$


(*delay coordinate AR-coefficients for stock*)
 Simplify[Coefficient[Expand[-Denominator[F[[5]]][[1]]], {e^{-i w}, e^{-2 i w}, e^{-3 i w}, e^{-4 i w}}]
 (*These are the s (t-1), ..., s (t-4) coefficients*)

$$\left\{ \frac{1-\beta}{1+\lambda+\lambda^2}, \frac{\lambda-\beta\lambda}{1+\lambda+\lambda^2}, -\frac{(-1+\beta)\lambda^2}{1+\lambda+\lambda^2}, 0 \right\}$$

(*AND delay coordinate MA-coefficients for stock*)
 Simplify[Coefficient[Expand[Numerator[F[[5]]][[1]]], {e^{-i w}, e^{-2 i w}, e^{-3 i w}, e^{-4 i w}}]
 (*These are the α (t), ..., α (t-3) coefficients*)

$$\left\{ \frac{1-\beta}{1+\lambda+\lambda^2}, \frac{\lambda-\beta\lambda}{1+\lambda+\lambda^2}, -\frac{(-1+\beta)\lambda^2}{1+\lambda+\lambda^2}, 0 \right\}$$

```
(*A PLOT OF NORMALIZED SPECTRAL DENSITIES:*)
 $\beta = .4$ ;  $\gamma = .3$ ;  $\lambda = 0.4$ ;
PSXN = PS[[1]][[1]] / NIntegrate[PS[[1]][[1]], {w, 0, Pi}];
PSYN = PS[[2]][[1]] / NIntegrate[PS[[2]][[1]], {w, 0, Pi}];
PSSN = PS[[5]][[1]] / NIntegrate[PS[[5]][[1]], {w, 0, Pi}];
p1 = Plot[{PSXN, PSYN, PSSN}, {w, 0, Pi},
  DefaultFont -> {"Times-New-Roman", 12},
  PlotStyle -> {Thickness[0.007],
    {Thickness[0.007], Dashing[{0.03, 0.03}]}, {Thickness[0.015]}},
  AxesStyle -> {Thickness[0.007]},
  AxesLabel -> {FontForm["Frequency", {"Times-New-Roman", 12}],
    FontForm["Power", {"Times-New-Roman", 12}]},
  Ticks -> {{0, 3}, {0, 1}},
  AspectRatio -> 1,
  PlotRange -> {0, 1.5}];
(* This is Fig 4 B in Bjornstad et al. -- Note though That the X-
  axis is in angular frequencies (0 -  $\pi$ ) *)
Clear[ $\beta$ ,  $\gamma$ ,  $\lambda$ ]
```



(*Calculating the limit of the Transfer functions as $d \rightarrow \text{Infinity}$

(cf. Table 2)*)

$d = \text{Infinity};$

(*defining the infinit denominators of the transfer functions*)

$\text{denominator} = (1 + \text{Exp}[-I w] \gamma -$

$(1 - \beta) \text{Exp}[-I w] (\text{Sum}[\lambda^{(n-1)} \text{Exp}[-n I w], \{n, 1, (d+1)\}] / \text{Sum}[\lambda^n, \{n, 0, d\}]));$

(* 0 gr transfer function in the limit as $d \rightarrow \text{Infinity}$ *)

$F0 = (1 + \text{Exp}[-I w] \gamma) / \text{denominator}$

$$\frac{1 + e^{-i w} \gamma}{1 + e^{-i w} \gamma - \frac{e^{-i w} (1-\beta) (1-\lambda)}{e^{i w - \lambda}}}$$

(* $k (> 0)$ gr transfer function in the limit as $d \rightarrow \text{Infinity}$ *)

$Fk = (1 - \beta) \text{Exp}[-I k w] / \text{denominator}$

$$\frac{e^{-i k w} (1 - \beta)}{1 + e^{-i w} \gamma - \frac{e^{-i w} (1-\beta) (1-\lambda)}{e^{i w - \lambda}}}$$

(* Stock transfer function in the limit as $d \rightarrow \text{Infinity}$ *)

$Fs = (\text{Exp}[-I w] (1 - \beta) (- \text{Sum}[\lambda^{(n-1)} \text{Exp}[-n I w], \{n, 1, (d+1)\}] / \text{Sum}[\lambda^n, \{n, 0, d\}])) /$
 denominator

$$- \frac{e^{-i w} (1 - \beta) (1 - \lambda)}{(1 + e^{-i w} \gamma - \frac{e^{-i w} (1-\beta) (1-\lambda)}{e^{i w - \lambda}}) (e^{i w} - \lambda)}$$

```
(*A PLOT OF NORMALIZED SPECTRAL DENSITIES
FOR THE LIMITING CASE: O-group and spawning stock*)
 $\beta = .4$ ;  $\gamma = .3$ ;  $\lambda = 0.4$ ;
T0 = ExpToTrig[Simplify[F0]];
TS = ExpToTrig[Simplify[Fs]];
PS0 = Re[T0]^2 + Im[T0]^2;
PSS = Re[TS]^2 + Im[TS]^2;
PS0N = PS0/NIntegrate[PS0, {w, 0, Pi}];
PSSN = PSS/NIntegrate[PSS, {w, 0, Pi}];
p1 = Plot[{PS0N, PSSN}, {w, 0, Pi},
  DefaultFont -> {"Times-New-Roman", 12},
  PlotStyle -> {Thickness[0.007],
    {Thickness[0.007], Dashing[{0.03, 0.03}]}, {Thickness[0.015]}},
  AxesStyle -> {Thickness[0.007]},
  AxesLabel -> {FontForm["Frequency", {"Times-New-Roman", 12}],
    FontForm["Power", {"Times-New-Roman", 12}]},
  Ticks -> {{0, 3}, {0, 1}},
  AspectRatio -> 1,
  PlotRange -> {0, 1.5}];
(* Note That the X-axis is in angular frequencies (0 -  $\pi$ ) *)
Clear[ $\beta$ ,  $\gamma$ ,  $\lambda$ ]
```

