



SPRINGTAILS

Order: Collembola

Springtails are very small, abundant, insect-like critters that inhabit a variety of moist environments. Although various springtail species range in size from 0.2 mm to 10 mm, most of the types that occasionally get into homes are 1- 2 mm in length. If they become a nuisance in homes, it is usually during extended droughts or prolonged, saturating rains.

DESCRIPTION

Springtails have antennae, three pairs of legs, a segmented body and can have simple mouthparts, such as piercing-sucking mouthparts, used to suck fluids from fungi or other liquid sources to very complex mouthparts that are highly specialized for some yet unknown reason. All of the different types of mouthparts are located and concealed within the head. It is principally this later characteristic that some authors believe separates the springtails from the rest of the class Insecta.


Their most identifiable morphological characteristic is called a furcula. It is a bi-forked appendage located on the underside of the fourth abdominal segment, is folded up underneath the springtail - held in place by a clasplike structure called the tenaculum - and is snapped downward to propel the springtail through the air. Presumably, this method of locomotion is used as a means to evade predators as it can propel the springtail up to 100 times its body length.

Springtails come in a variety of colors and shapes. Most are elongated, but some species are more globular. Their colors range from white to black, and some species are purple, red, orange, grey, yellow or mottled (multicolored).

LIFE HISTORY AND BEHAVIOR

Springtails require high humidity environments such as beneath leaf litter, in mulch, under thatch in damp lawns similar locations. Their numbers have been estimated to be as high as 100,000 per cubic meter (approximately 35 cubic feet). They feed on molds, mildew, fungi, bacteria and decaying plant material, occasionally causing damage to crops in gardens, fields, greenhouses, and to cultivated mushrooms.

In homes with high humidity and mold growth, springtails can sometimes maintain an indoor population. This is more common in bathrooms and basements but can also occur within walls of



Several types of Springtails found in structures.

newly built homes that are supporting mold growth because of rains that soaked the floor and walls before they were covered. One species of springtails, *Hypogastrura nivicola*, are known as snow fleas. On warm winter days these black springtails can sometimes be seen on the snow surface searching for pollen spores. This event is more common in or adjacent to wooded areas.

MANAGEMENT

During droughty conditions, springtails will move from mulches and landscaping in search of more humidity. As previously mentioned, water saturated soils will also drive springtails from their habitats in search of, in this case, drier conditions. As the springtails seek more favorable environs they can inadvertently enter homes under doors, sill plates, siding and windows, and through screens.

The best method to manage or eliminate springtails in the home is by lowering the humidity through the use of air conditioning or de-humidifiers. This is of particular relevance in bathrooms, showers and saunas. Also, any mold and mildew should be removed.

Water house plants thoroughly and then allow the soil to dry before watering again.

Reduce the amount of mulch surrounding the home.

Hire a pest management professional (pest control company) to apply an insecticide to the exterior foundation and building perimeter.

WARNING

Pesticides are poisonous. Read and follow directions and safety precautions on labels. Handle carefully and store in original labeled containers out of the reach of children, pets, and livestock. Dispose of empty containers right away, in a safe manner and place. Do not contaminate forage, streams, or ponds.

Steven B. Jacobs
Sr. Extension Associate
Dept. of Entomology
March 2010

HP-26

©The Pennsylvania State University 2010

This publication is available in alternative media on request.

Where trade names are used, no discrimination is intended and no endorsement by The Pennsylvania State University or Pennsylvania Department of Agriculture is implied.

Entomological Notes are intended to serve as a quick reference guide and should not be used as a substitute for product label information. Although every attempt is made to produce Entomological Notes that are complete, timely, and accurate, the pesticide user bears the responsibility of consulting the pesticide label and adhering to those directions.

Issued in furtherance of Cooperative Extension Works, Acts of Congress May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture and the Pennsylvania Legislature. D. Jackson, Director of Cooperative Extension, The Pennsylvania State University.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. It is the policy of the University to maintain an academic and work environment free of discrimination, including harassment. The Pennsylvania State University prohibits discrimination and harassment against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, or veteran status. Discrimination or harassment against faculty, staff, or students will not be tolerated at The Pennsylvania State University. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 328 Bouke Building, University Park, PA 16802-5901, Tel 814-865-4700/V, 814-863-1150/TTY.