



## EASTERN TENT CATERPILLAR

*Malacosoma americanum* (Fabricius)

The eastern tent caterpillar has been observed in the United States since 1646. Outbreaks frequently occur at eight to ten year intervals. The presence of this pest in the spring is usually recognized by their conspicuous nests or tents constructed in the forks and crotches of a tree. Before gypsy moth outbreaks in the 1970s, some experts considered this pest to be a significant defoliator of deciduous shade trees in the northeastern United States.

### DESCRIPTION

The egg mass (Fig. 1a) of this species encircles small twigs and appears to be varnished. The egg mass may be nineteen millimeters. Small larvae spin fine strands of silk wherever they crawl. As the larvae grow, so does the size of the tent. Fully grown larvae are about two inches long, generally black with a white stripe down the middle of the back. Mature caterpillars (Fig. 1b) will leave the host tree to search for a suitable place to spin their pale yellowish cocoons. During late June and July the reddish-brown adult moths with two oblique, white bands on the forewing emerge from their cocoons (Fig. 1c). After mating, the female deposits eggs in a mass around small twigs on a host plant.

### LIFE HISTORY

This key pest overwinters as a dark, glistening, collar-like egg mass on twigs. Each egg mass contains 150 to 350 eggs. After hatching from eggs in the spring, about the time cherry leaves issue from buds, the gregarious larvae start to construct their tents in nearby branch crotches. These tents are made of silken layers and serve as protective sites for larvae. The caterpillars feed for a period of six to eight weeks. The mature caterpillars have a habit of migrating to protected areas to spin their cocoons. Within the cocoon the larva transforms into a resting stage called the pupa. The pupal stage lasts about three weeks. During late June and July adults emerge and the female deposits eggs in a mass around small twigs. There is only one generation a year in Pennsylvania.




Figure 1. Eastern tent caterpillar viable egg mass (a), mature larva (b), and adult (c).

### DAMAGE

The preferred hosts of this pest are cherry, crabapple, and apple. The eastern tent caterpillar occasionally attacks other deciduous ornamental shrubs, shade, and forest trees. The silky tents spun by the caterpillars make landscape trees unsightly, and the caterpillars are annoying when searching for food or a suitable place to spin their cocoons.

The caterpillars do not feed within their webs, but congregate there during the night and rainy weather. When caterpillars are abundant, they frequently eat all the leaves on a tree which weakens it, but seldom kills it. The foliage on the host tree may be stripped from all the twigs within a distance of three feet from the nest(s).

## MANAGEMENT

### **(Only for non-edible ornamental trees and shrubs)**

#### **Non-Chemical**

Eastern tent caterpillar larvae are prey for other insects, toads, and birds. Several kinds of small, beneficial wasps parasitize eggs, larvae, or pupae of this pest. Many caterpillars die from disease during unfavorable weather. Prune small twigs that have viable egg masses on them from December through March before egg hatch.

#### **Chemical**

Registered insecticides should be applied in April when the caterpillars are small. To control the eastern tent caterpillars, apply registered formulations to the nests and about one foot of the surrounding branches or trunk. Apply the spray before the nests are three inches in diameter. If possible, make applications early in the morning when the caterpillars are generally inside their nests. Follow all label directions for specific information on host plant label clearance, phytotoxicity information, safety precautions, and dosage information.

## WARNING

Pesticides are poisonous. Read and follow directions and safety precautions on labels. Handle carefully and store in original labeled containers out of the reach of children, pets, and livestock. Dispose of empty containers right away, in a safe manner and place. Do not contaminate forage, streams, or ponds.

Gregory A. Hoover  
Sr. Extension Associate  
Dept. of Entomology  
March 2002

TS-12

©The Pennsylvania State University 2002

This publication is available in alternative media on request.

Where trade names are used, no discrimination is intended and no endorsement by The Pennsylvania State University or Pennsylvania Department of Agriculture is implied.

Entomological Notes are intended to serve as a quick reference guide and should not be used as a substitute for product label information. Although every attempt is made to produce Entomological Notes that are complete, timely, and accurate, the pesticide user bears the responsibility of consulting the pesticide label and adhering to those directions.

Issued in furtherance of Cooperative Extension Works, Acts of Congress May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture and the Pennsylvania Legislature. T.R. Alter, Director of Cooperative Extension, The Pennsylvania State University.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. It is the policy of the University to maintain an academic and work environment free of discrimination, including harassment. The Pennsylvania State University prohibits discrimination and harassment against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, or veteran status. Discrimination or harassment against faculty, staff, or students will not be tolerated at The Pennsylvania State University. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 201 Willard Building, University Park, PA 16802-2801, Tel 814-865-4700/V, 814-863-1150/TTY.