



# Entomological Notes

Department of Entomology

## EUROPEAN HORNET

*Vespa crabro* L.

The European or giant hornet is an introduced species first reported in the United States in 1840 in New York. Currently, its geographical range extends from the Northeastern states west to the Dakotas, and south to Louisiana and Florida. It belongs to a family of wasps called the Vespids, which encompass all of the yellowjackets including the bald-faced hornets. Technically, it is the only hornet in North America. The European hornet is a large, aggressive yellowjacket. Homeowners should be cautious when attempting to control their nests.

### DESCRIPTION

The adult European hornet worker is approximately 35 mm in length with yellow and brown coloration (Fig. 1). The overwintering queens are somewhat larger. The nests are typically located in a cavity, such as a hollow tree or wall void. They will rarely appear freely suspended like the football-shaped bald-faced hornet nests. The entrance to European hornets' nests are frequently two meters (six feet) or more aboveground. In some instances, a portion of the gray, papery nest extends outside the cavity or void.

### LIFE HISTORY

Each fall, the colony produces males and females that mate, and the females become next year's queens. Only the overwintering queens survive in protected sites such as under loose bark, in tree cavities, and in wall voids of buildings. All other colony members produced in the current year will perish.

In the spring, the emerging queens establish new nests in aerial cavities, deposit eggs in cells they have constructed, and feed the first batch of larvae. The larvae mature, pupate in their cells, and then emerge as sterile female workers. These workers take over the responsibility of foraging for food to feed the young larvae, collect cellulose to expand the nest, and protect the nest from external threats. Typical food for the young include crickets, grasshoppers, large flies, caterpillars, and the workers of other yellowjacket species.




Figure 1. European hornet

European hornet colonies often contain 300 or more workers by September or October (maximum of 800–1,000 workers). These workers are unique among the yellowjackets for their ability (actually a propensity) to forage at night. It is not unusual for workers to bounce off of external lights or house windowpanes during summer nights. Although the workers will sting if handled, they are not normally aggressive unless the colony is threatened.

In addition to the hazard created by their stings, the hornets will also damage various trees and shrubs by girdling the branches and twigs to gather bark for nest building and to obtain nourishment from the sap.

### MANAGEMENT

For treatment of European hornets in wall voids of buildings, we advise the use of professional pest control services. Be certain NOT to plug the hornets' entrance because they may chew through interior wall coverings in an attempt to escape and enter the living area.

### WARNING

Pesticides are poisonous. Read and follow directions and safety precautions on labels. Handle carefully and store in original labeled containers out of the reach of children, pets, and livestock. Dispose of empty containers right away, in a safe manner and place. Do not contaminate forage, streams, or ponds.

Steven B. Jacobs  
Sr. Extension Associate  
Dept. of Entomology  
March 2000  
Reviewed September 2003

PH-7

© The Pennsylvania State University 2003

This publication is available in alternative media on request.

Where trade names are used, no discrimination is intended and no endorsement by The Pennsylvania State University or Pennsylvania Department of Agriculture is implied.

Entomological Notes are intended to serve as a quick reference guide and should not be used as a substitute for product label information. Although every attempt is made to produce Entomological Notes that are complete, timely, and accurate, the pesticide user bears the responsibility of consulting the pesticide label and adhering to those directions.

Issued in furtherance of Cooperative Extension Works, Acts of Congress May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture and the Pennsylvania Legislature. T.R. Alter, Director of Cooperative Extension, The Pennsylvania State University.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. It is the policy of the University to maintain an academic and work environment free of discrimination, including harassment. The Pennsylvania State University prohibits discrimination and harassment against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, or veteran status. Discrimination or harassment against faculty, staff, or students will not be tolerated at The Pennsylvania State University. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 328 Bouke Building, University Park, PA 16802-5901, Tel 814-865-4700/V, 814-863-1150/TTY.