



Corrigendum: Coupling Developmental Physiology, Photoperiod, and Temperature to Model Phenology and Dynamics of an Invasive Heteropteran, *Halyomorpha halys*

Anne L. Nielsen^{1*}, Shi Chen² and Shelby J. Fleischer³

¹ Department of Entomology, Rutgers University, Bridgeton, NJ, United States, ² Department of Population Health and Pathobiology, North Carolina State University, Charlotte, NC, United States, ³ Department of Entomology, Pennsylvania State University, University Park, PA, United States

Keywords: brown marmorated stink bug, phenology, agent-based model, stochastic model, life-history, population dynamics, invasive species

A corrigendum on

Coupling Developmental Physiology, Photoperiod, and Temperature to Model Phenology and Dynamics of an Invasive Heteropteran, *Halyomorpha halys*

by Nielsen, A. L., Chen, S., and Fleischer, S. J. (2016). *Front. Physiol.* 7:165. doi: 10.3389/fphys.2016.00165

OPEN ACCESS

Edited and reviewed by:

Petros Damos,
Aristotle University of Thessaloniki,
Greece

*Correspondence:

Anne L. Nielsen
nielsen@aesop.rutgers.edu

Specialty section:

This article was submitted to
Invertebrate Physiology,
a section of the journal
Frontiers in Physiology

Received: 08 June 2017

Accepted: 20 July 2017

Published: 11 August 2017

Citation:

Nielsen AL, Chen S and Fleischer SJ
(2017) Corrigendum: Coupling
Developmental Physiology,
Photoperiod, and Temperature to
Model Phenology and Dynamics of an
Invasive Heteropteran, *Halyomorpha*
halys. *Front. Physiol.* 8:568.
doi: 10.3389/fphys.2017.00568

In this manuscript, we present model results from eight locations over 10 years based on temperatures at weather stations and photoperiod. We inadvertently made a poor choice for a weather station to represent Wenatchee WA. To avoid heat island effects, we chose the Grouse Camp weather station to represent Wenatchee. However, although Grouse Camp is only 21 km from Wenatchee, it is in a mountainous area (1,642 m elevation) and poorly represents the climate in the apple growing region of Wenatchee, WA (190 m elevation). Thus, the results do not reflect potential population dynamics of *H. halys* in Wenatchee. In re-evaluation of the model, the data show that populations at Wenatchee, WA, are predicted to behave similarly to those at Salem, OR, with an average albeit marginal positive population growth. Conclusions that were driven heavily by photoperiod, such as the range in days for initiation of oviposition by overwintered adults, were less affected (from 16 down to 13 days). The strong differences were due to markedly higher degree day accumulations at Wenatchee versus Grouse Camp.

The following six files use the same order of tables and figures from the original manuscript and use the Weather Station data for Wenatchee, WA (network ID: GHCND:USC00459074).

- **Table 2.** Model outputs defining key population parameters for the years 2005–2014.
- **Figure 7.** Model predictions of adult population size for Wenatchee, WA. *P* represented parental overwintered adults, which was initialized as 1000 for each year and for each simulation run.
- **Figure 8.** Predicted population sizes (+/– range from all simulations and years) across all geographic locations for (A) maximum adult population size, and (B) final population size, and (C) yearly degree-day accumulation. The error bars represent the standard errors from 100 simulations for the metrics.

- Figure S3. Predicted total population size by life stage for Wenatchee, WA from 2005 through 2014. Populations were initialized with 1000 adults for each year and simulation run.
- Figure S5. Degree-day accumulation for *Halyomorpha halys* development in Wenatchee, WA from 2005 through 2014.
- Figure S7. Predicted total population size by generation for Wenatchee, WA.

REFERENCES

Weather Station: Wenatchee, WA (Network ID: GHCND:USC00459074).

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

We thank V. Jones for bringing this error to our attention, and the Frontiers journal for allowing us to add this Corrigendum.

ACKNOWLEDGMENTS

This material is based upon the work supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number SCRI 2011-51181-30937.

Copyright © 2017 Nielsen, Chen and Fleischer. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 2 | Model outputs defining key population parameters for the years 2005-2014.

Location	Coordinates	Crop	Non-diapause range	P Oviposition		F ₁ Eclosion		F ₂ Eclosion		Final adult population	
				Range	Median	Range	Median	Range	Median	F ₁	F ₂
Geneva NY	42.88°N 76.99°W	Apple	Apr 18–Aug 26	Jun 7	Jun 21–Jul 18	Jun 30	Jul 19–Aug 9	Jul 28	243–447	81–1,847	
Bridgeton NJ	39.43°N 75.23°W	Peach/Vegetable	Apr 22–Aug 22	Jun 4	Jun 9–Jul 2	Jun 23	Jul 6–Jul 31	Jul 24	140–278	531–2,027	
Asheville NC	35.58°N 85.56°W	Tree fruit/Vegetable	Apr 28–Aug 17	May 29	Jun 3–Jun 11	Jun 6	Jun 25–Jul 4	Jul 2	89–253	803–2,287	
Homestead FL	25.47°N 80.47°W	Tomato/Strawberry	May 24–July 22	Jun 6	Jun 17–Jun 18	Jun 18	Jul 8–Jul 10	Jul 9	44–137	1318–2,781	
Wenatchee WA	47.42°N 120.33°W	Apple/Pear	Apr 14–Aug 31	Jun 8	Jun 17–Jun 29	Jun 22	Aug 1–Aug 18	Aug 6	106–559	46–1,557	
Salem OR	44.93°N 123.03°W	Tree fruit/Wine grape	Apr 17–Aug 28	Jun 9	Jun 17–Jul 2	Jun 26	Jul 24–Jul 31	Jul 27	269–716	33–1,035	
Davis CA	38.55°N 121.74°W	Tomato	Apr 24–Aug 20	Jun 4	Jun 14–Jun 19	Jun 16	Jul 6–Jul 15	Jul 11	75–255	733–1,893	
Riverside CA	33.95°N 117.40°W	Citrus	May 1–Aug 13	Jun 11	Jun 20–Jun 24	Jun 23	Jul 8–Jul 16	Jul 13	21–95	349–962	




FIGURE 7 | Model predictions of adult population size for Wenatchee, WA. *P* represented parental overwintered adults, which was initialized as 1,000 for each year and for each simulation run.

