

Seasonal Forcing on the Dynamics of *Clethrionomys rufocanus*: Modeling Geographic Gradients in Population Dynamics

Nils Chr. STENSETH^{*,†,1)}, Ottar N. BJØRNSTAD^{*,2)} and Takashi SAITOH^{†,‡,3)}

Researches on Population Ecology (1998): **40**:85-94

* *Division of Zoology, Department of Biology, University of Oslo, P.O. Box 1050 Blindern, N-0316 Oslo, Norway.* † *Center of Advanced Studies, Drammensvein 78, N-0271, Oslo, Norway.* ‡ *Hokkaido Research Center, Forestry and Forest Products Research Institute, Sapporo 062-8516, Japan*

1) E-mail: n.c.stenseth@bio.uio.no

2) E-mail: o.n.bjornstad@bio.uio.no

3) E-mail: bedford@ffpri-hkd.affrc.go.jp

Abstract. We interpret the gradient in population dynamics from the southwestern part of the island of Hokkaido to its northeastern part within the framework of a phenomenological model. In Hokkaido, as in other northern regions, both spring and fall is considered as short transition periods between the two main seasons – summer (the primary breeding season) and winter (the non-reproductive or secondary breeding season). We show that the geographic transition in dynamics may be understood as the combined consequence of different patterns of density-dependence during summer and winter, and geographically varying season lengths. Differences are shown to exist between summer and winter with respect to strength of density-dependence. Direct density-dependence, in particular, is stronger during winter than during summer. A model is presented to show how relative lengths of seasons can induce both stable and periodically fluctuating population dynamics. The results are compared and contrasted with what is otherwise known about the gradient in rodent dynamics in Fennoscandia.

Key words: density-dependence, length of summer and winter, phenomenological model, seasonal and annual dynamics, the grey-sided vole.

Introduction

Populations of the grey-sided vole (*Clethrionomys rufocanus* (Sundevall, 1846)) in the southern and western parts of Hokkaido exhibit only seasonal dynamics with population growth during summer and decline during winter. Populations in northern and eastern parts of Hokkaido exhibit multiannual periodic density fluctuations in addition to the seasonal dynamics (Bjørnstad et al. 1998a; Saitoh et al. 1998a; see also Saitoh 1987; Bjørnstad et al. 1996;

Stenseth et al. 1996a; Saitoh et al. 1998b). The cycles in these northeastern parts resemble those found in microtine rodents in Fennoscandia (e.g. Hansson and Henttonen 1988; Hanski et al. 1991, 1993; Stenseth and Ims 1993; Bjørnstad et al. 1995; Stenseth et al. 1996b). The regional transition from pure seasonal to multiannually fluctuations is associated with an increase in population variability and a transition in the patterns of direct and delayed density-dependence (Stenseth et al. 1996a; Saitoh et al. 1998a).

Various mechanisms have been suggested to explain the biogeographic transitions in dynamics of small rodents: changes in community composition (notably the predator guild; Erlinge 1987; Hansson and Henttonen 1988; Hanski et al. 1991; Bjørnstad et al. 1995; Turchin and Hanski 1997), changes in primary production (Jedrzejewski and Jedrzejewska 1996), and changes in climatic conditions (notably seasonality and extent of snowcover; Hansson and Henttonen 1985; Hansson 1987; Bjørnstad et al. 1998a). In Hokkaido, the main correlate to this transition appears to be the relative length of the summer (Kaneko et al. 1998). The most obvious difference between the seasons is with respect to reproduction (winter generally being a non-reproductive season; see, e.g. Ota 1984; Kaneko et al. 1998). A few recent studies on microtine rodents have, however, also indicated that patterns of density-dependence may differ between seasons (Mallorie and Flowerdew 1994; Hansen et al. 1998).

In this paper we explore the hypothesis that the relative lengths of seasons are responsible for the transition in dynamics. We propose that density-dependence differs between summer and winter, so that the overall strengths of direct and delayed density-dependence are shifted with the relative lengths of the seasons, which again affect the dynamical properties of the populations. For the purpose of providing background information, we first briefly summarize the difference in the growing season between southwestern and northeastern Hokkaido, and indicate how patterns of density-dependence differ between winter and summer. We then develop a phenomenological model that allows us to investigate the effect of relative lengths of summer and winter on annual patterns of density-dependence. Throughout we adopt the approach of Stenseth et al. (1996b) by asking what model (and mechanism) is compatible with the observed patterns. We close the paper with a general discussion with respect to microtine rodent dynamics and with respect to dynamical effects of seasonal forcing.

Seasonal patterns of density-dependence

The climate in Hokkaido is affected by latitude and altitude as well as the direction of sea currents (see Kaneko et al. 1998). Warm currents run along the




Fig. 1. (A) The geographic location of the 11 groups of *Clethrionomys rufocanus* populations, and (B) the relation between the 'index of vegetation growth period' (discussed in this paper) and the geographic score index (representing the 'northeasterlyness' and discussed by Saitoh et al. (1998a)). The line in B represents the linear regression; the Pearson correlation coefficient is -0.71 ($P = 0.01$). The 225 time series discussed and analyzed in Saitoh et al. (1998a) were, as described by Saitoh et al. (1998a), grouped into the 11 groups according to geographic and climatic criteria: Group 1 (Teshio) consists of 31 time series; Group 2 (Kitami) consists of 31 time series; Group 3 (Abashiri) consists of 20 time series; Group 4 (Yubari) consists of 8 time series; Group 5 (Kamikawa-Taisetsu) consists of 27 time series; Group 6 (Shiribeshi) consists of 27 time series; Group 7 (West-Hidaka) consists of 20 time series; Group 8 (Oshima) consists of 30 time series; Group 9 (East-Hidaka) consists of 10 time series; Group 10 (Tokachi) consists of 13 time series; Group 11 (Kushiro-Nemuro) consists of 8 time series. The length of the time series are (depending on group) between 12 and 31 year (194 are longer than 23 years).

western and southern shores of Hokkaido resulting in a milder climate for these regions (see Saitoh et al. 1998a). A northern cold current prevails along the northern and eastern shores resulting in a cooler climate for this region. The growing season for plants varies accordingly. In the southwest the growing season is long whereas in the northeast it is shorter (with a corresponding longer winter). An 'index of vegetative growth period' (also referred to as a

WI) for the growing season of plants was presented and discussed by Kira (1949, 1971); it is defined as the sum average of monthly temperatures above 5°C within a year. Thus, $WI = \sum (T - 5)$, where T is average monthly temperature (excluding months of which the average temperature is lower than 5°C). The larger the value of the index, the longer is the growing season. This index correlates well with the forest distribution of Japan (Kira 1949, 1971). The 'geographic score index' (see Saitoh et al. 1998a) which correlates with the transition in dynamics from the southwestern to the northeastern part of Hokkaido (see Saitoh et al. 1998a) also correlates crudely with the 'index of vegetative growth period' (*WI*). We use the *WI*-index as a proxy for length of the breeding period of voles. This period is known to be around 7.5 months in the southwestern part of Hokkaido and around 6 month in the northeast (Fujimaki 1975; Kaneko et al. 1998).

The population dynamics of *C. rufocanus* has been studied on the basis of 225 time series on rodent abundance from the census conducted by the Forestry Agency of the Japanese Government (FAJ) (Saitoh et al. 1998a; see also Kaneko et al. 1998; Bjørnstad et al. 1998a). (Most of the time series are between 20 and 31 years long; see Kaneko et al. (1998).) Populations were sampled (as described by Kaneko et al. (1998) in both spring (May/June) and fall (September/October) as well as in the middle of the summer (July/August). Saitoh et al. (1998a) used the fall samples to estimate the annual structure of density-dependence in the 225 time series. The fall-samples are from the end of the breeding season (cf. Kaneko et al. 1998); hence, between two subsequent fall-samplings there is only one continuous sequence of winter (non-reproductive season) and summer (reproductive season).

For characterizing the population growth from spring to fall and from fall to spring, we use in this paper also the spring samples.

The index of vegetative growing period (*WI*) is not yet available for all the 225 individual time series from the FAJ. As discussed by Saitoh et al. (1998a), the 225 time series may be grouped into 11 groups, according to geography and topographical characteristics (Fig. 1A). In this paper we adopt the regions depicted in Fig. 1A as our basic units. Figure




Fig. 2. The autoregressive coefficients (see Eq. (1)) estimated from the average time series from each group (see Fig. 1A; see also Bjørnstad et al. 1998a: Fig. 1) plotted against the ‘index of the vegetation growth period’. (A) The 1st order coefficient ($R^2 = 0.50$, $P = 0.012$). (B) The 2nd order coefficient ($R^2 = 0.10$, $P = 0.35$).

1B depicts the relation between the index of vegetative growth period (WI) and the geographic score, which is based on the location order on the two directions (from south to north and from west to east) to each group (see also Saitoh et al. 1998a). The autoregressive analysis reported below is primarily based on the combined (i.e. mean) time series within each of the 11 groups (see Bjørnstad et al. 1998a: Fig. 1). However, when analyzing winter and summer density-dependence (see *Discussion*) we use data from the individual census points from the northern part of Hokkaido (roughly corresponding to the data used by Stenseth et al. (1996a)).

Saitoh et al. (1998a) demonstrate that several population dynamical descriptors correlate with the geographic score. The strength of density-dependence is, in particular, found to change with geographic position (Saitoh et al. 1998a; see also Stenseth et al. 1996a). This pattern remains when focusing on the descriptors of each of the mean time series for each group (see Bjørnstad et al. 1998a: Fig. 1), although less clearly so.

We measure strength of direct and delayed density-dependence using the second order autoregressive model applied to the log-transformed series of fall abundance (Royama 1992; Bjørnstad et al. 1995):

$$x_t = b_0 + b_1 x_{t-1} + b_2 x_{t-2} + e_t \quad (1)$$

where x_t is the log-transformed fall abundance (abundance + 1) of year t , b_1 and b_2 are the first and second order autoregressive coefficients, b_0 is a scaling parameter, and $\{e_t\}$ is a sequence of independent and identically distributed random variables with zero mean and a constant variance (see also Stenseth et al. 1996a,b).

The first order autoregressive coefficient (b_1) increases with the index for growing period (Fig. 2A). There is a less clear relationship between the second order autoregressive coefficient (b_2) and the index for the growing season (Fig. 2B). (Stenseth et al. (1996a) reported a slightly different gradient. This is, however, due to the fact that only the northern subset of the database was analyzed then since only this part of the data was available at the time. Closer inspection show that the gradient observed by Stenseth et al. (1996a) may be seen as a part of the overall gradient when the entire island is considered.) Altogether it appears that the ‘index of vegetative WI ’ is a better correlate than the ‘geographic scores’; unfortunately the ‘index of vegetative growth period’ is not yet available for all individual time series where the ‘geographic scores’ are.

In the following section we explore the hypothesis that the geographic gradient in the density-dependent structure of the time series may be understood as a result – direct or indirect – of relative season length. The present exposition should be considered a preliminary report only. The pattern of seasonal differences will certainly need more careful study. We hope that the present exposition will stimulate further work on the issue of seasonal differences in density-dependence and the resulting dynamic effects.

A model of seasonal dynamics

The log-linear autoregressive model: a strategic preamble

The exact mechanism causing the microtine cycle is still in debate (see, e.g. Stenseth et al. 1998a). It is

clear, though, that the mechanisms responsible for these periodic fluctuations lead to a rather specific signature of density-dependence (Stenseth et al 1996a,b, 1998b; see also Bjørnstad et al. 1995). The second order autoregressive log-linear model (Eq. (1)) has been found to be appropriate for the fall time series on *C. rufocanus* in northern Hokkaido (Stenseth et al. 1996a). A similar model structure has been successfully fitted to other microtine populations (see Bjørnstad et al. 1995; Stenseth et al. 1996b). The linear autoregressive model can arise (to some level of approximation) from a variety of more specific ecological (e.g. trophic) annual models, and can be interpreted in terms of such (see Stenseth et al. 1996a, 1997 for detailed examples). The general seasonal mechanism we propose does, however, not hinge on any specific ecological model. We therefore use the second order model as a starting point, and leave the issue of the exact mechanisms behind the density-dependence to be discussed elsewhere.

The estimated coefficients for the mean time series for each of the 11 groups are shown in Figs. 2 and 3. The distinct geographic pattern observed in the level of variability and periodicity (Bjørnstad et al. 1998a; Saitoh et al. 1998a) is reflected in the autoregressive coefficients. The gradient in the first order coefficient is very clear (Figs. 2 and 3). The patterns in the second order coefficient is less clear-cut, but the western groups (1, 6, 8) stand out by not exhibiting delayed density-dependence (Fig. 3). On the basis of the second order autoregressive model, northeast areas are expected to exhibit from 3.5 to 4.5 year cycles (Fig. 3B; see also Bjørnstad et al. 1995).

Incorporating seasonality

Assume N_t to be the fall-abundance of year t resulting from population growth during the preceding year, we may write the following model for the skeleton (i.e. the deterministic component of the dynamics):

$$N_t = N_{t-1} \exp(a_0 - a_1 x_{t-1} - a_2 x_{t-2}), \quad (2)$$

where $\exp(a_0)$ is the annual maximum per capita multiplication rate, a_1 measures the reduction in this growth rate due to direct density-dependence and a_2 that due to delayed density-dependence (see, e.g. Stenseth et al. 1996a). Note that positive values for the parameters a_1 and a_2 represents negative density-dependence.

To divide the annual growth into a component due to winter dynamics and another due to summer dynamics, it is easy to see that the model (2) may be split in a multiplicative manner (see Åström et al. 1996). Let t ($0 < t < 1$) be the relative length of the summer defined as lasting from the initiation of spring reproduction until the end of the reproduction

in fall. The annual model may then be seen as a sequence of density-dependent growths according to:

$$N_t = N_{t-1} \exp[(a_{w0} - a_{w1} x_{t-1} - a_{w2} x_{t-2})(1-t)] \exp[(a_{s0} - a_{s1} x_{t-1} - a_{s2} x_{t-2})t], \quad (3)$$

where the parameters with the w -subscript refer to winter (expressed as per- t units) and the parameters with the s -subscript refer to summer. During the summer two demographic processes operate (survival and reproduction), whereas during the winter only one process operates (survival; winter reproduction is generally assumed to be negligible for *C. rufocanus* in Hokkaido; see Ota 1984; Kaneko et al. 1998; Yoccoz et al. 1998).

Log-transforming both sides of Eq. (3) and collecting terms, we may write:

$$x_t = a_{w0}(1-t) + a_{s0}t + (1-a_{w1}(1-t) - a_{s1}t)x_{t-1} - (a_{w2}(1-t) + a_{s2}t)x_{t-2}. \quad (4)$$

The parameters in the model given by Eq. (4) may be directly related to the components of the autoregressive model given by Eq. (1):

$$b_0 = a_{w0}(1-t) + a_{s0}t, \quad (5a)$$

$$b_1 = 1 - a_{w1} + (a_{w1} - a_{s1})t, \quad (5b)$$

$$b_2 = -a_{w2} + (a_{w2} - a_{s2})t. \quad (5c)$$

Changes in the length of the breeding season (t) can thus potentially induce a transition in the dynamics if the patterns of density-dependence differ between the seasons (below).

The data on Clethrionomys rufocanus

We may compare equation (5) with the relations depicted in Fig. 2. Since the first order autoregressive coefficient increases with the length of the summer period, t , we need to have $(a_{w1} - a_{s1}) > 0$ implying $a_{w1} > a_{s1}$. That is, the intraspecific interactions during the winter (measured per t -units) is stronger than during the summer.

To expand on this result and help interpret the implication of changing the relative length of the summer (t), we have investigated a specific numerical example (Fig. 4). The length of the breeding (summer) period (for which the WI -index has been used as a proxy in this paper) in the southwestern and the northeastern part of Hokkaido corresponds to $t = 0.625$ and $t = 0.5$, respectively (see above). Consider a line in the autoregressive parameter space (Fig. 4A; defined by the regression lines shown in Fig. 2) connecting the longest and the shortest growth period (long: $b_1 = 0.02$ and $b_2 = -0.26$; short: $b_1 = -0.24$ and $b_2 = -0.31$). Assume that all constants in equation (3) are positive (i.e. we only allow for negative density-dependence). We can then solve for the unknown parameters in equation (5), which gives: $a_{w1} = 1.14$, $a_{w2} = 0.651$, $a_{s1} = 0.2$ and a_{s2}




Fig. 3. (A) Map of the autoregressive coefficients for the time series of each of the 11 groups. The first column represent the 1st order coefficient, the second column represents the 2nd order coefficient. Positive values are shaded in grey. (B) A scatter plot of the coefficients for the 11 groups. The predicted dynamics based on the second order autoregressive model are given by the contour lines (see Royama 1992; Bjørnstad et al. 1995). Outside the triangle, the dynamics is divergent. Below the semicircle, dynamics is cyclic. The contour lines give the periodicity (measured in years) of the cycle. The arrow connects the groups with the highest WI to that of the lowest WI among the 11 groups.

= 0. The conclusion is clear: density-dependence (as measured per time unit; e.g. per t -units) is profoundly stronger during the winter than during the summer.

Using the season-specific parameters, the gradient resulting from $t = 0.5$ to 0.63 translates to the arrow shown in Fig. 4A. (The full transition from pure winter ($t = 0$) to pure summer dynamics ($t = 1$) is given by the darker line). The resulting gradients in population dynamics and periodic length are shown in Fig. 4B. The model spans the range of observed periodicities (see Bjørnstad et al. 1998a); however, the gradient from $t = 0.5$ to 0.63 fails to predict the exact transition from stable to cyclic dynamics.

If we, instead consider the full range in parameters observed in the data (Fig. 3) and relax the assumption that all parameters in equation (3) must be positive (which may be realistic, at least with respect to delayed density-dependence), we obtain $a_{w1} = 2.40$, $a_{w2} = 0.651$, $a_{s1} = 0$ and $a_{s2} = -0.35$. (Again the density-dependence during the winter is stronger than during the summer.) The resultant gradient from the southwestern to the northeastern part of Hokkaido is represented by the arrow in Fig. 4C. (The full transition from pure winter ($t = 0$) to pure summer dynamics ($t = 1$) is given by the darker line.) This refined model predicts that southeastern populations (where the summer is long, $t = 0.63$) should have no multiannual periodicity (i.e. only exhibit seasonal fluctuations) and that northeastern populations (where the summer is short, $t = 0.5$) have a 3-4 year cycle. The model is also able to capture the rather surprising effect that longer cyclic periods are observed at intermediate lengths of summer (between the stable and the 3-4 year cycle), as is the case for the central

part of Hokkaido (Group 4; see Bjørnstad et al. 1998a: Fig. 2).

The pattern of seasonal differences will certainly need future and more careful scrutiny. The present preliminary results do, however, demonstrate the viability of our basic hypothesis, namely that the relative lengths of the reproductive season may generate – directly or indirectly – the observed geographic pattern in population dynamics of *C. rufocanus* in Hokkaido. The intriguing finding is that we need not assume any particular clinal changes in the underlying density-dependent parameters (measured per time unit) to explain the transition in dynamics. It suffices to assume that the seasonal driver shifts the relative importance of the winter and summer components of the population dynamics.

Discussion

We have focused on whether seasonal differences in density-dependence in combination with large-scaled geographic differences in season lengths may explain the shift in dynamics from purely seasonal dynamics in the south and west to the multiannual cycles in the northeast of Hokkaido. For this to be so, density-dependence during winter needs to be stronger than during summer. Altogether our analysis demonstrates that seasonality (or seasonal forcing) may represent a key factor to population cycles in small rodents, which originally was suggested by Saitoh (1987). There are several theoretical studies on how seasonal forcing can act as an agent to induce population cycles (e.g. Kot and Schaffer 1984; Rinaldi et al. 1993; Grenfell et al. 1995). These studies show that seasonality is a potent bifurcation agent, if it expresses itself as cyclic variation in one (or more) of the model parameters. The current study approaches




Fig. 4. Numerical examples to illustrate how changing the relative length of the summer (τ) changes the dynamics of a population when density-dependence differs between the seasons (see Eq. (4)). (A and B) We assume that the patterns of summer and winter density-dependence are $a_{w1} = 1.14$, $a_{w2} = 0.651$, $a_{s1} = 0.2$ and $a_{s2} = 0$. (A) Annual patterns of density dependence as a function of season length (the line spans $t = 0$ to $t = 1$). The arrow covers the distance from $t = 0.63$ to $t = 0.75$. (B) The resulting period length as a function of t . A period length of zero corresponds to noncyclic dynamics. As can be seen, in a region with mainly summer, there is no cycles whereas cycles of length 3-4 years are expected in regions with short summers. Notice that after a short transition range, the period length is fairly insensitive to the relative length of the seasons. (C and D) The same as A and B but with $a_{w1} = 2.40$, $a_{w2} = 0.651$, $a_{s1} = 0$ and $a_{s2} = -0.35$.

this general issue from a related but more phenomenological angle by focusing on seasonality in terms of what Åström et al. (1996) coined sequential density-dependence. Åström et al. (1996) speculated that sequential density-dependence models may be useful for understanding cyclic rodent dynamics; we concur with this.

In the following discussion we will first focus the seasonal difference, and provide some preliminary analyses showing that density-dependence is indeed stronger during winter. Then, we discuss our findings in the more general context of gradients in rodent dynamics.

Seasonal differences in density-dependence

In order to explore the validity of our major conclusion of the stronger density-dependence during the winter, we have undertaken an analysis to investigate differences in density-dependence between summer and winter in the grey-sided vole populations of Hokkaido. We used 73 time series from Northern Hokkaido that have complete spring

and winter censuses for all the 31 years of study (1963-1993). These results are preliminary, firstly, because some assumptions need to be made, and secondly because some methodological studies remain to be undertaken. The spring and autumn censuses are roughly 6 months apart. By assuming that the reproductive season in Northern Hokkaido is around 6 months (i.e. $t \approx 0.5$; see above) we may assess a_{w1} and a_{s1} from the census data (a_{w2} and a_{s2} may also be assessed; but for our preliminary exposition, a_{w1} and a_{s1} will suffice). In the absence of measurement error, estimates of a_{w1} and a_{s1} can be obtained through regressing $\log(\text{spring abundance})$ on $\log(\text{preceding autumn abundance})$ and *vice versa* (see Royama 1992). However, the rodent censuses are a count variable which does not provide exact estimates of abundance. Count variables typically harbour a Poisson or negative binomial error (e.g. Pielou 1977; Lebreton 1990). Measurement error bias estimates of density-dependence upwards (the density-dependence is concluded to be stronger than it really is) when using the 'log-regression' method (e.g. Itô 1972; see Bjørnstad et al. (1998b) for details when the error is of the Poisson-type). The key




Fig. 5. Scatterplot of the estimates of winter density-dependence against summer density-dependence for each 73 time series from Northern Hokkaido that have complete spring and winter censuses for all the 31 years of study (1963-1993). The estimates assume that the error in the dependent census count follows a Poisson error (see text). The basic model is: $Nf_t = Ns_t \exp[c_s - b_{summer} \ln Ns_t]$, where Ns_t and Nf_t are spring and fall abundances (not log-transformed), b_{summer} is the density-dependence over the summer; c_s is the density-independent summer growth rate. The basic model being assumed for the winter growth is: $Ns_t = Nf_{t-1} \exp[c_w - b_{winter} \ln Nf_{t-1}]$, where b_{winter} is the density-dependence over the winter; c_w is the density-independent winter growth rate. Notice that the length of the winter and the summer in the particular region from which the data derive are about equal to each other ($b_{winter} \approx a_{s,t} 0.5$ and $b_{winter} \approx a_{w,t} 0.5$). The straight line represents identical strengths of density-dependence in the winter and in the summer. The scatter suggests that density-dependence is strongest in the winter. But a *provisio* must be made about the measurement error (see text).

methodological problem for seasonal comparisons is that the variance of counts depends on the mean. For estimates based on annual counts (as those reported in Fig. 2), this bias is present: but it will be of the same magnitude for most populations. However, in seasonal estimation the bias may well differ between winter estimates (for which the initial abundance is relatively high) and summer estimates (for which the initial abundance is relatively low). In our calculations we estimate density-dependence assuming the dependent count has a Poisson error (e.g. Lebreton 1990; Bjørnstad et al. 1998b). We are in the process of developing methods that accomodates the Poisson error in the independent counts (using Markov chain Monte Carlo methods; e.g. Gilks et al. 1996). Figure 5 depicts the winter and summer estimates of density-dependence for each of the 73 populations. The winter estimates are invariably greater than the summer estimates. With a *provisio* for any artefacts induced by differences in the magnitude of measurement error, we thus have

some indications that winter density-dependence is stronger than summer density-dependence in these populations. Altogether then, these results support our basic assertion that density-dependence differs between seasons, and that it is strong during winter.

Mechanisms for seasonal differences

Our discussion on mechanisms behind the stronger winter density-dependence must necessarily remain speculative. The conclusion that winter regulation is the strongest is, in fact, surprising for a number of reasons: Reproduction is for one thing known to be density-dependent in these voles (e.g. Saitoh et al. 1997; Kaneko et al. 1998). Spacing behavior should, if anything, be expected to lead to stronger intraspecific interactions during the summer, since females defend exclusive territories during the breeding season, but have overlapping home ranges in winter (Ishibashi et al. 1998a,b). Food may be limiting in winter (Petrušewicz et al. 1971; Bujalska 1975; Viitala 1977). However, *C. rufocanus* of Hokkaido eat leaves and shoots of bamboograsses (*Sasa kurilensis*, *S. senanensis*, *S. nipponica* and others) for foods in winter (Ota 1984). Bamboograss is the most common undergrowth in forests of Hokkaido and it has never been reported that bamboograss is depleted by vole's grazing. Thus, bamboograss seems super-abundant for the voles of Hokkaido, though its nutritious quality may not be high in winter (which after all may lead to resource limitations). Cockburn (1988: Chapt. 5) reviews differences in survival, reproduction, and social behavior between spring-born and autumn-born cohorts. It may be worth to revisit some of these differences to explain the patterns of direct density-dependence. For now, however, the pattern remains enigmatic: it seems real but unexpected.

Any difference in delayed density-dependence, if present, is as that for direct density-dependence. Even though there are many possible scenarios, the argument can be made (cf. Stenseth et al. 1996b) that delayed density-dependence often seems to be induced by the trophic interaction between voles and some specialized predators such as mustelids. The abundance of rodents in the proceeding year, can then be seen as a proxy for the abundance of specialist predators in the year in question. Hansson (1987) suggested that the interaction between rodents and their predator may be modified greatly by snow cover (see also Lindström and Hörnfeldt 1994; Saitoh and Takahashi 1998). If predation is the source of delayed density-dependence in rodents, any such effect could lead to seasonal differences. This is another important topic for future study, since delayed density-dependence is a critical component of the rodent cycles (Hörnfeldt 1994; Bjørnstad et al. 1995; Stenseth et al. 1996a,b).




Fig. 6. (A) The map of the autoregressive coefficients for time series of various microtine rodents at 10 different locations in Fennoscandia (data as analyzed in Stenseth et al. 1996b: Table 1). When more than one species are present, the coefficients represent the averages. The first column represent the 1st order coefficient, the second column represents the 2nd order coefficient. Positive values are shade in grey. (B) A scatter plot of the coefficients for the 10 locations. The predicted dynamics are given by the contour lines (see Fig. 4). The arrow connects the southernmost and the northernmost location. The species, location and sources are as follows: 1) *Lemmus lemmus* and *Microtus oeconomus* at Finse from Framstad et al. (1993, 1997); 2) *Clethrionomys rufocanus* at Boda from Marcström et al. (1990); 3, 4) Both *Microtus* spp at Alajoki and Routsala from Korpimäki and Norrdahl (1991), Korpimäki (1993, 1994); 5) *C. glareolus* and *M. agrestis* at Sotkamo from Henttonen et al. (1977), A. Kaikusalo (personal communication); 6) *C. rufocanus*, *C. glareolus* and *M. agrestis* at Umeå Hörnfeldt (1994); 7) *C. rufocanus*, *C. glareolus* and *C. rutilus* at Pallasjärvi from Henttonen et al. (1977, 1987), Hanski and Henttonen (1996), H. Henttonen (personal communication); 8) *C. rufocanus* on Kola from Kataev et al. (1994); 9) *C. glareolus* on Kola from Koshkina (1966) (provided by P. Turchin); and 10) *C. rufocanus* at Kilpisjärvi from Kalela (1957), Laine and Henttonen (1983, 1987), H. Henttonen (personal communication).

Gradients in rodent dynamics

To complement the discussion of the dynamics in Hokkaido, we briefly revisit the dynamics of microtines in Fennoscandia (Hansson and Henttonen 1988; Hanski et al. 1991; Bjørnstad et al. 1995). In Fennoscandia there are pure seasonal fluctuations in the south and multiannually periodic fluctuations in the north (see Hansson and Henttonen 1988). In addition, there is a clear latitudinal gradient in patterns of statistical density-dependence. Bjørnstad et al. (1995) showed that the latitudinal gradient in the periodicity of the cyclic rodents of Fennoscandian (i.e. those north of 60°N) is associated with a decrease in the strength of direct density-dependence (Fig. 6). A dominant hypothesis in the literature to explain this gradient is through changes in the predator community (Hanski et al. 1991; Bjørnstad et al. 1995; Turchin and Hanski 1997). Leaving aside the cyclicity itself, the analysis presented in this paper suggests an alternative explanation for the gradient north of approximately 60°N, namely that a shift in season length may change the annual pattern of density-dependence (see also Hansen et al. 1998). This may be a particularly fruitful line of inquiry for Fennoscandian rodents, since the length of the winter changes greatly with latitude. Figure 6 depicts the

gradient in the autoregressive coefficients: in the southern part the first autoregressive coefficient is negative whereas in the northern part it is positive. This is in stark contrast to the pattern of correlation between density-dependence and season length presented for Hokkaido. One possible difference could be that subnivean winter reproduction appears to be quite common in parts of Fennoscandia (cf. Stenseth 1978; Hansson 1984). The seasonal differences with respect to parameter values seem to be different from the Hokkaido case: within the framework of our model (Eq. (2)), we would need to assume that $a_{w1} < a_{s1}$ (incidentally, this may be consistent with assuming that generalist predators play an important role in Fennoscandia but hunting success differs between seasons; cf. Hanski et al. 1991)¹.

This is not the place to enter into a detailed analysis of the Fennoscandian gradient. The above considerations do, however, demonstrate that our model (and hypothesis) may profitably be applied to

¹ Realizing that in Kilpisjärvi (northern Finland), the summer is very short (i.e. t is small) and the winter is long (i.e. $1-t$ is large), we might easily obtain $a_{w1}(1-t) > a_{s1}t$ (as concluded by Hansen et al. 1998).

other rodent populations. It is a rather general phenomenological hypothesis pointing to the importance of the relative length of the seasons, as well as detailed characteristics of the conditions during the different seasons.

Conclusion

The most important general conclusion of our analysis is that seasonal forcing and seasonal differences between density-dependencies may represent a key to the understanding of microtine population dynamics. Within the field of microtine rodent ecology, there has been a disproportional focus on the summer season even though pioneers in the field (e.g. Elton 1924; Formozov 1946; Chitty 1952; Pitelka 1957; Kalela 1962; Schwarz 1963, 1977; Schwarz et al. 1964; Fuller 1967; Merritt 1984) all emphasized the importance of winter regulation and abrupt seasonality. In regions where multiannual fluctuations are most common, the winter covers a substantial portion of the year. It is our hope that this paper will contribute to a renewed focus on seasonality and the winter ecology of rodents.

Acknowledgements: Support from Forestry and Forest Products Research Institute of Japan for bringing NCS to Japan (1998) is greatly appreciated. This paper was conceived during his staying in Japan (and on travel back to Europe). A grant from Japan Society for Promotion Science, given to H. Abe for bringing NCS to Japan (in 1991), made it possible to start the collaboration between Norway and Japan for population ecology of the grey-sided vole; a follow-up grant from the Norwegian Research Council, given to NCS for bringing TS to Norway (in 1994 and 1997) facilitated our studies. Support from the Nansen Endowment for bringing ONB to Japan (in 1996) is greatly appreciated. The Center for Advanced Study (Oslo, Norway) also supported this project (in 1997, during which some focus was devoted to the effect of seasonality). Many discussions with Thomas Hansen on the issue of seasonality in dynamics of periodically fluctuating populations has influenced us profoundly and is greatly appreciated. Comments by Hirofumi Hirakawa, Romain Julliard, Mauricio Lima, Anne-Caroline Prévot-Julliard and Nigel G. Yoccoz on an earlier draft of the paper are greatly appreciated.

References

- Åström, M., P. Lundberg and S. Lundberg (1996) Population dynamics with sequential density-dependencies. *Oikos* **75**: 174-181.
- Bjørnstad, O. N., W. Falck and N. C. Stenseth (1995) A geographic gradient in small rodent density fluctuations: a statistical modelling approach. *Proceedings of the Royal Society of London B* **262**: 127-133.
- Bjørnstad, O. N., S. Champely, N. C. Stenseth and T. Saitoh (1996) Cyclicity and stability of grey-sided voles, *Clethrionomys rufocanus*, of Hokkaido: spectral and Principal components analyses. *Philosophical Transactions of the Royal Society of London Series B* **351**: 867-875.
- Bjørnstad, O. N., N. C. Stenseth, T. Saitoh and O. C. Lindgjærde (1998a) Mapping the regional transition to cyclicity in *Clethrionomys rufocanus*: spectral densities and functional data analysis. *Researches on Population Ecology* **40**: ###-###.
- Bjørnstad, O. N., J.-M. Fromentin, N. C. Stenseth and J. Gjøsaeter (1998b) A test for density-dependent survival in juvenile stages of cod. (in review)
- Bujalska, G. (1975) Reproduction and mortality of bank voles and the changes in the size of an island population. *Acta Theriologica* **20**: 41-56.
- Chitty, D. (1952) Mortality among voles (*Microtus agrestis*) at Lake Vyrnwy, Montgomeryshire in 1936-39. *Philosophical Transactions of the Royal Society of London Series B* **236**: 505-552.
- Cockburn, A. (1988) *Social behaviour in fluctuating populations*. Croom Helm.
- Elton, C. S. (1924) Periodic fluctuations in the numbers of animals: their causes and effects. *Journal of Experimental Biology* **2**: 119-163.
- Erlinge, S. (1987) Predation and noncyclicity in a microtine population in Southern Sweden. *Oikos* **50**: 347-352.
- Formozov, A. N. (1946) Snow cover as an integral factor of the environment and its importance in the ecology of mammals and birds. *Materials Fauna and Flora of USSR, New Series, Zoology* **5**: 1-152 (originally in Russian).
- Framstad, E., N. C. Stenseth and E. Østbye (1993) Time series analysis of population fluctuations of *Lemmus lemmus*. pp. 97-115. In N. C. Stenseth and R. A. Ims (eds.) *The biology of lemmings*. Academic Press, London.
- Framstad, E., N. C. Stenseth, O. N. Bjørnstad and W. Falck (1997) Limit cycles in Norwegian Lemmings: tensions between phase-dependence and density-dependence. *Proceedings of the Royal Society of London B* **264**: 31-38.
- Fujimaki, Y. (1975) Reproductive activity in *Clethrionomys rufocanus bedfordiae* 3. Regional differences in autumn reproductive activity. *Bulletin of the Hokkaido Forest Experiment Station* **13**: 38-45. (in Japanese with English summary)
- Fuller, W. A. (1967) Ecologie hivernale des lemmings et fluctuations de leurs populations. *Terre Vie* **114**: 97-115.
- Gilks, W. R., S. Richardson and D. J. Spiegelhalter (eds.) (1996) *Markov chain Monte Carlo in practice*. Chapman & Hall, London.

- Grenfell, B. T., B. M. Bolker and A. Kleczkowski (1995) Seasonality and extinction in chaotic metapopulations. *Proceedings of the Royal Society of London B* **259**: 97-103.
- Hansen, T., N. C. Stenseth and H. Henttonen (1998) Multiannual vole cycles and population regulation during long winters: an analysis of seasonal density dependence. (in review)
- Hanski, I. and H. Henttonen (1996) Predation on competing rodent species: a simple explanation of complex patterns. *Journal of Animal Ecology* **65**: 220-232.
- Hanski, I., L. Hansson and H. Henttonen (1991) Specialist predators, generalist predators, and the microtine rodent cycle. *Journal of Animal Ecology* **60**: 353-367.
- Hanski, I., P. Turchin, E. Korpimäki and H. Henttonen (1993) Population oscillations of boreal rodents: Regulation by mustelid predators leads to chaos. *Nature* **364**: 232-235.
- Hansson, L. (1984) Winter reproduction of small mammals in relation to food conditions and population dynamics. pp. 225-234. In J. F. Merritt (ed.) *Winter ecology of small mammals*. Special Publication of Carnegie Museum of Natural History, 10. Pittsburgh.
- Hansson, L. (1987) An interpretation of rodent dynamics as due to trophic interactions. *Oikos* **50**: 308-318.
- Hansson, L. and H. Henttonen (1985) Gradients in density variations of small rodents: the importance of latitude snow cover. *Oecologia* **67**: 394-402.
- Hansson, L. and H. Henttonen (1988) Rodent dynamics as community processes. *Trends in Ecology and Evolution* **3**: 195-200.
- Henttonen, H., A. Kaikusalo, J. Tast and J. Viitala (1977) Interspecific competition between small rodents in subarctic and boreal ecosystems. *Oikos* **29**: 581-590.
- Henttonen, H., T. Oksanen, A. Jortikka and V. Haukialmi (1987) How much do weasels shape microtine cycles in the northern Fennoscandian taiga? *Oikos* **50**: 353-365.
- Hörfeldt, B. (1994) Delayed density dependence as a determinant of vole cycles. *Ecology* **75**: 791-806.
- Ishibashi, Y., T. Saitoh and M. Kawata (1998a) Social organization of the vole *Clethrionomys rufocanus* and its demographic and genetic consequences: a review. *Researches on Population Ecology* **40**: ###-###.
- Ishibashi, Y., T. Saitoh, S. Abe and M. C. Yoshida (1998b) Kin-related social organization in a wintering population of the vole *Clethrionomys rufocanus*. *Researches on Population Ecology* **40**: ###-###.
- Itô, Y. (1972) On the methods of determining density-dependence by means of regression. *Oecologia* **10**: 347-372.
- Jedrzejewski, W. and B. Jedrzejewska (1996) Rodent cycles in relation to biomass and productivity of ground vegetation and predation in the Palearctic. *Acta Theriologica* **41**: 1-34.
- Kalela, O. (1957) Regulation of reproduction rate in subarctic populations of the vole, *Clethrionomys rufocanus* (Sund.). *Annales Academiae Scientiarum Fennicae, Series A, IV Biologica* **34**: 1-60.
- Kalela, O. (1962) On the fluctuations in the numbers of arctic and boreal small rodents as a problem of production biology. *Annales Academiae Scientiarum Fennicae, Series A, IV, Biologica* **66**: 1-38.
- Kaneko, Y., K. Nakata, T. Saitoh, N. C. Stenseth and O. N. Bjørnstad (1998) The biology of the vole *Clethrionomys rufocanus*: a review. *Researches on Population Ecology* **40**: ###-###.
- Kataev, G. D., J. Suomela and P. Palokangas (1994) Densities of microtine rodents along a pollution gradient from a copper-nickel smelter. *Oecologia* **97**: 491-498.
- Kira, T. (1949) Fundamental problems of climate classification. *Seibutsu Kagaku* (Biological Science) **1**: 193-199. (in Japanese)
- Kira, T. (1971) *Nature from the ecological view point*. Kawaide-shobo, Tokyo. (in Japanese)
- Korpimäki, E. (1993) Regulation of multiannual vole cycles by density-dependent avian and mammalian predation. *Oikos* **66**: 359-363.
- Korpimäki, E. (1994) Rapid or delayed tracking of multi-annual vole cycles by avian predators? *Journal of Animal Ecology* **63**: 619-628.
- Korpimäki, E. and K. Norrdahl (1991) Do breeding nomadic avian predators dampen population fluctuations of small mammals? *Oikos* **62**: 195-208.
- Koshkina, T. V. (1966) On the periodical changes in the numbers of voles. *Bulletin of the Moscow Society of Naturalists, Biological Section* **71**: 1-26.
- Kot, M. and W. M. Scahffer (1984) The effects of seasonality on discrete models of population growth. *Theoretical Population Biology* **26**: 340-360.
- Laine, K. and H. Henttonen (1983) The role of plant production in microtine cycles in northern Fennoscandia. *Oikos* **40**: 407-418.
- Laine, K. and H. Henttonen (1987) Phenolics/nitrogen ratio in the bilberry *Vaccinium myrtillus* in relation to temperature and microtine density in Finnish Lapland. *Oikos* **50**: 389-395.
- Lebreton, J.-D. (1990) Modelling density dependence, environmental variability, and demographic stochasticity from population counts: An example using Wytham Wood great tits. pp. 89-102. In J. Blondel, A. Gosler, J.-D. Lebreton and R. McCleery (eds.) *Population biology of the passerine*

- birds. Springer-Verlag, Berlin.
- Lindström, E. R. and B. Hörnfeldt (1994) Vole cycles, snow depth and fox predation. *Oikos* **70**:156-160.
- Marcström, V., N. Höglund and C. J. Krebs (1990) Periodic fluctuations in small mammals at Boda, Sweden from 1961 to 1988. *Journal of Animal Ecology* **59**: 753-761.
- Merritt, J. F. (ed.) (1984) *Winter ecology of small mammals*. Special Publication of Carnegie Museum of Natural History, 10. Pittsburgh.
- Mallorie, H. C. and J. R. Flowerdew (1994) Woodland small mammal population ecology in Britain: a preliminary review of the Mammal Society survey of wood mice *Apodemus sylvaticus* and bank voles *Clethrionomys glareolus*, 1982-87. *Mammal Review* **24**: 1-15.
- Ota, K. (ed.) (1984) *Study on wild murid rodents in Hokkaido*. Hokkaido University Press, Sapporo. (in Japanese)
- Petrusewicz, K., G. Bujalska, R. Andrzejewski and J. Gliwicz (1971) Productivity processes in an island population of *Clethrionomys glareolus*. *Annales Zoologici Fennici* **8**: 127-132.
- Pielou, E. C. (1977) *Mathematical ecology*. Wiley & Sons, New York.
- Pitelka, F. A. (1957) Some aspects of population structure in the short-term cycle of the brown lemming in Northern Alaska. *Cold Spring Harbor Symposia on Quantitative Biology* **18**: 237-251.
- Rinaldi, S., S. Muratori and Y. Kuznetsov (1993) Multiple attractors, catastrophes and chaos in seasonally perturbed predator prey communities. *Bulletin of Mathematical Biology* **55**: 15-35.
- Royama, T. (1992) *Analytical population dynamics*. Chapman and Hall, London.
- Saitoh, T. (1987) A time series and geographical analysis of population dynamics of the red-backed vole in Hokkaido, Japan. *Oecologia* **73**: 382-388.
- Saitoh, T. and K. Takahashi (1998) The role of vole populations in prevalence of the parasite (*Echinococcus multilocularis*) in foxes. *Researches on Population Ecology* **40**: ###-###.
- Saitoh, T., N. C. Stenseth and O. N. Bjørnstad (1997) Density dependence in fluctuating grey-sided vole populations. *Journal of Animal Ecology* **66**: 14-24.
- Saitoh, T., N. C. Stenseth and O. N. Bjørnstad (1998a) The population dynamics of the vole *Clethrionomys rufocanus* in Hokkaido, Japan. *Researches on Population Ecology* **40**: ###-###.
- Saitoh, T., O. N. Bjørnstad and N. C. Stenseth (1998b) Density-dependence in voles and mice: a comparative study. *Ecology* (in press).
- Schwarz, S. S. (1963) Adaptations of terrestrial vertebrates to the environmental conditions of the subarctic. Academy of Sci. USSR, Ural Affiliat, Inst. Biol., Trudy **33**: 1-19 (originally in Russian).
- Schwarz, S. S. (1977) *The evolutionary ecology of animals*. Consultant Bureau, New York. (originally in Russian)
- Schwarz, S. S., A. V. Pokrowski, V. G. Istchenko, V. G. Olenjey, N. A. Ovtschinnikova and O. A. Pjastolova (1964) Biological peculiarities of seasonal generation of rodents with special references to the problem of senescence in mammals. *Acta Theriologica* **8**: 11-43.
- Stenseth, N. C. (1978) Demographic strategies in fluctuating populations of small rodents. *Oecologia* **33**: 149-172.
- Stenseth, N. C. and R. A. Ims (eds.) (1993) *The Biology of lemming*. Academic Press, London.
- Stenseth, N. C., O. N. Bjørnstad and T. Saitoh (1996a) A gradient from stable to cyclic populations of *Clethrionomys rufocanus* in Hokkaido, Japan. *Proceedings of the Royal Society of London B* **263**: 1117-1126.
- Stenseth, N. C., O. N. Bjørnstad and W. Falck (1996b) Is spacing behaviour coupled with predation causing the microtine density cycle? A synthesis of process-oriented and pattern-oriented studies. *Proceedings of the Royal Society of London B* **263**: 1423-1435.
- Stenseth, N. C., W. Falck, O. N. Bjørnstad and C. J. Krebs (1997) Population regulation in snowshoe hare and lynx populations: asymmetric food web configurations between the snowshoe hare and the lynx. *Proceedings of the National Academy of Science for the USA* **94**: 5147-5152.
- Stenseth, N. C., T. Saitoh and N. G. Yoccoz (1998a) Frontiers in population ecology of microtine rodents: a pluralistic approach to the study of population ecology. *Researches on Population Ecology* **40**: ###-###.
- Stenseth, N. C. R. Boonstra, N. G. Yoccoz and C. J. Krebs (1998b) Population cycles in lemmings and voles: dynamics and demographic signatures of non-linearity. (in review)
- Turchin, P. and I. Hanski (1997) An empirically based model for latitudinal gradient in vole population dynamics. *American Naturalist* **149**: 842-874.
- Viitala, J. (1977) Social organization in cyclic subarctic populations of the voles *Clethrionomys rufocanus* (Sund.) and *Microtus agrestis* (L.). *Annales Zoologici Fennici* **14**: 53-93.
- Yoccoz, N. G., K. Nakata, N. C. Stenseth and T. Saitoh (1998) The demography of *Clethrionomys rufocanus*: from mathematical and statistical models to further field studies. *Researches on Population Ecology* **40**: #####-#####.