

Antennal Lobe Partitioning of Behaviorally Active Odors in Female Cabbage Looper Moths

J. L. Todd, T. C. Baker

Department of Entomology, Iowa State University, Ames, Iowa, 50011-3222, USA

A highly specific cobalt marking method for moth peripheral receptors [1] has led to great progress in characterizing the functional organization of the glomeruli within the male-specific macroglomerular complex (MGC) by focusing on first-order processing of female-emitted sex pheromones [1–4]. The first evidence for anatomically and functionally distinct glomeruli within the MGC was shown with *Manduca sexta* L. in a study of projection patterns of pheromone-sensitive interneurons [5]. The pathways of sensory axons into antennal lobe glomeruli of female moths have thus far been examined in only one noctuid species, *Spodoptera littoralis* (Boisd.), and these neurons were tuned to the conspecific major sex pheromone component [3]. The glomerular destinations of axons from sensory neurons specific for other behaviorally active odor blend components besides female-emitted sex pheromones have not heretofore been investigated for any moth species.

The main objective of the present study was to examine first-order processing of floral odors and male- and female-produced pheromone components in female cabbage looper moths, *Trichoplusia ni* (Hübner). Both the male courtship pheromone (one component of which is linalool) and an odor bouquet from flowers of *Abelia grandiflora* (phenylacetaldehyde, 2-phenylethanol, benzyl alcohol, benzaldehyde) have been shown to evoke upwind flight in *T.*

ni females [6, 7], and single-neuronal recordings have revealed neurons that are tuned to individual components of these behaviorally active odor blends [8]. Female *T. ni* also have antennal neurons that exhibit a low threshold similar to that of the male's neurons [9] for the major component of the female-emitted sex pheromone, (*Z*)-7-dodecenyl acetate (*Z*7-12:Ac) [9], although the behavioral significance for this detection is not known. Because upwind flight in female *T. ni* can be caused by floral odors [6] or by the male courtship pheromone [7], we became interested in determining whether neurons tuned to components of these two odor blends project into overlapping or separate areas of the antennal lobe, thus stimulating upwind flight by activating the same olfactory pathways and perhaps the same sensation of odor, or by using different pathways that result in a unique type of odor quality integration, respectively. Rearing and maintenance of *T. ni* females is according to [8]. Descriptions of the cut-sensillum extracellular recording technique [10], cobalt marking method, and neuroanatomical reconstruction of the antennal lobes and axonal pathways follow in full the methodology described for male *T. ni* [4]. The odor stimuli used and their preparation and presentation follow that of [8]. The sustained pulsing of a sex pheromone component preferentially stains the neuron within the cut sensillum that is

tuned to the pulsed component [1–4]; the specificity of the staining for other odorant-tuned neurons has not been investigated prior to this study.

The morphology of female *T. ni* antennal lobes was determined from 10- μ m frontal sections, and measurements of the anterior-to-posterior (longitudinal) dimensions of 38 antennal lobes (Table 1) indicated that glomerular tissue spans $249.1 \pm 27.8 \mu\text{m}$. Female moth brains have thus far reflected varied glomerular architecture [3, 12–14]. There was considerable variation in size among the individual glomeruli making up each lobe of female *T. ni*. The largest glomeruli were not consistently located at the entrance of the antennal nerve into the antennal lobe, as in male moths [1–4], in which two complexes of glomeruli are clearly present [1–5, 11]. Only a few females had one or more enlarged glomeruli, or a complex of ordinary-sized glomeruli, located at the dorsalmost aspect of the lobe, that were clearly separated from the more ventrally situated glomeruli. There was no identifiable, consistent arrangement of glomeruli having the appearance of the *T. ni* MGC [4] in conspecific females, nor did the ordinary glomeruli appear in a consistent location or with reproducible shapes from individual to individual based on light-microscopic examination of serial sections. Willis et al. [15] also found no stereotyped arrays of identifiable glomeruli in the antennal lobes of female *M. sexta*. In contrast, Rospars' [16] quantitative morphometric analysis of the glomerular organization of the antennal lobes of *Mamestra brassicae* indicated that the shape, size, and relative positions of glomeruli in both sexes were markedly constant, such that each glomerulus could be assigned a number and be recognized either within or between individual moths.

We connected successfully with 341 trichoid sensilla on female *T. ni* antennae, as indicated by spontaneous background activity; however, the receptor neurons within only 71 of these sensilla were excited by one or more of the six odorants that we tested. Spontaneous firing of neurons within most of the sensilla that we sampled indicated at least three different spike amplitudes. Todd and Baker [8] have previously illustrated simultaneously recorded action potentials and DC potentials of female *T. ni* receptor neurons in response to the same set of stimuli used in the present study.

We successfully stained neurons with cobalt in 38 of the 71 sensilla wherein spike activity was recorded such that axonal projections could be traced into the antennal lobe (Table 1). In 31 of the 38 sensilla, a single receptor neuron was stained (single stain), and in seven sensilla, two neurons were stained (double stain; Table 1). In all of the single stains, the axon terminated within a single glomerulus. Regardless of whether neurons were tuned to one of the four floral compounds or to the male- or female-emitted pheromone component, they projected their axons into glomeruli located at the dorsal-most aspect of the lobe, closest to the antennal nerve entrance. No lateral or medial location ("addresses") were consistently targeted by neurons tuned to a particular component or to a type of odor (floral, courtship pheromone, sex pheromone). Thus, in female *T. ni*, it appears that the dorsalmost glomeruli in the antennal lobe process a relatively diverse array of odor signals, many of

which are known to evoke upwind flight. The corresponding area in male moth antennal lobes is thus far exclusively devoted to input concerning the female-emitted sex pheromone [1-4] or to the breakdown product of the major sex pheromone component of the blend [1, 4], all of which are known to modulate upwind flight. The processing role of glomeruli in the central and ventral regions of the antennal lobe of female *T. ni* was not illuminated by using our six odorant stimuli.

Serial sections revealed that there does appear to be a longitudinal partitioning of axonal terminations from neurons tuned to the components that make up different odor blends (Table 1). Axons from neurons tuned to floral odorants tended to terminate in glomeruli located at the posterior of the antennal lobe, whereas projections from neurons transmitting information about the female-emitted sex pheromone component were located in glomeruli positioned closest to the anterior aspect of the lobe, as indicated by the center-point of each of the arborizations in the serial sections (Table 1). This pattern of partitioning was also evident when the size of the individual female's antennal lobe was taken into account (Table 1, right column). Glomeruli involved in processing information about the courtship pheromone were located more medially within the glomerular sac (Table 1).

There is a growing body of evidence that morphologically distinct glomeruli of the MGC [1-4], or even subunits of a single glomerulus [4], represent functionally distinct processing centers for

incoming information about individual sex pheromone components; however, no such spatially consistent functional separation into single glomeruli seems to exist in female *T. ni* for individual floral compounds or for sex-pheromone or courtship-pheromone components. This does not mean that there might not be several groups of functionally specific glomeruli that collectively receive input from the same exclusive type of neuron as suggested by the longitudinal partitioning data above. Thus far, we have been unable to identify consistent and unique spatial locations of individual functionally distinct glomeruli in female *T. ni*.

We thank E. W. Meador for rearing the moths, and D. Sakaguchi for use of equipment necessary for facilitating antennal lobe reconstructions. This research was supported by NSF grant No. 93-09948 to TCB. Journal paper No. J-16578 of the Iowa Agriculture and Home Economics Experiment Station, Ames, Iowa, Project No. 3240, and supported by Hatch Act and State of Iowa funds.

Received October 17, 1995 and March 22, 1996

1. Hansson, B.S., et al.: Science 256, 1313 (1992)
2. Hansson, B.S., et al.: J. Comp. Physiol. A 177, 535 (1995)
3. Ochieng', S.A., et al.: Tissue Cell 27, 221 (1995)
4. Todd, J.L., et al.: Physiol. Entomol. 20, 349 (1995)
5. Hansson, B.S., et al.: J. Comp. Neurol. 312, 264 (1991)

Table 1. Antennal lobe partitioning of axonal projections of odorant-specific sensory neurons in female *T. ni* trichoid sensilla

Stimulus	Attempted stains	Successful stains		Mean projection depth ^a in antennal lobe [$\mu\text{m} \pm \text{SD}$]	Front-to-back length of projection [μm]	Relative projection depth per antennal lobe depth
		Single	Double			
Benzaldehyde	7	4	0	142.5 \pm 24.7	15.0 \pm 7.1	0.67 \pm 0.1
Benzyl alcohol	14	9	1	187.8 \pm 26.6	40.0 \pm 15.8	0.78 \pm 0.9
2-Phenylethanol	15	5	2	177.1 \pm 45.4	42.9 \pm 17.9	0.74 \pm 0.2
Phenylacetaldehyde	14	5	0	166.3 \pm 50.6	27.5 \pm 9.6	0.61 \pm 0.2
Linalool	10	2	2	125.0 \pm 42.4	20.0 \pm 14.1	0.53 \pm 0.2
Z7-12:Ac	11	6	2	86.3 \pm 13.1	22.5 \pm 15.0	0.37 \pm 0.1
Total attempts	71					
Total stains		31	7			

^a Numbers increasing in size represent depths of glomerular tissue located toward the posterior of the antennal lobe; measurements based on single stains only

6. Haynes, K. F., et al.: *J. Chem. Ecol.* 17, 637 (1991)
 7. Landolt, P. J., Heath, R. R.: *Science* 249, 1026 (1990)
 8. Todd, J. L., Baker, T. C.: *Naturwissenschaften* 80, 183 (1993)
 9. Todd, J. L., et al.: *Physiol. Entomol.* 17, 183 (1992)
 10. Kaissling, K.-E., in: *Biochemistry of Sensory Functions*, p. 243 (L. Jaenicke, ed.). Berlin: Springer 1974
 11. Matsumoto, S. G., Hildebrand, J. G.: *Proc. R. Soc. London Ser. B* 213, 249 (1981)
 12. Christensen, T. C., Hildebrand, J. G., in: *Functions, Organization, and Physiology of the Olfactory Pathways in the Lepidopteran Brain*, p. 457 (A. P. Gupta, ed.). New York: Wiley 1987
 13. Anton, S., Hansson, B. S.: *J. Comp. Physiol. A* 350, 199 (1994)
 14. Koontz, M. A., Schneider, D.: *Cell Tissue Res.* 249, 39 (1987)
 15. Willis, M. A., et al.: *J. Comp. Physiol. A* 176, 205 (1995)
 16. Rospars, J. P.: *J. Comp. Neurol.* 220, 80 (1983)
-