

The Biology of the Vole *Clethrionomys rufocanus*: a Review

Yukibumi Kaneko, Keisuke Nakata, Takashi Saitoh, Nils Chr. Stenseth and Ottar N. Bjørnstad

Researches on Population Ecology (1998): 40:21-37

Biological Laboratory, Faculty of Education, Kagawa University, Takamatsu 760-0016 Japan. Hokkaido Forestry Research Institute, Bibai 079-0198, Japan. Hokkaido Research Center, Forestry and Forest Products Research Institute, Sapporo 062-8516, Japan. Division of Zoology, Department of Biology, University of Oslo, P.O. Box 1050 Blindern, N-0316 Oslo, Norway

1) E-mail: kaneko@laplace.ed.kagawa-u.ac.jp

2) E-mail: nakata@hfri.bibai.hokkaido.jp

3) E-mail: bedford@ffpri-hkd.affrc.go.jp

4) E-mail: n.c.stenseth@bio.uio.no

5) E-mail: o.n.bjornstad@bio.uio.no

Abstract. The biology of the grey-sided vole *Clethrionomys rufocanus* in Hokkaido, concerning taxonomy, morphology, phylogeny, distribution, and natural history, is reviewed. Applied issues in forest management (damage, control and census) are also historically mentioned. Although *Clethrionomys rufocanus* of Hokkaido was originally identified as a distinct species, *Evotomys* (= now *Clethrionomys*) *bedfordiae* Thomas, 1905, current literature generally refers to the grey-sided vole of Hokkaido as *Clethrionomys rufocanus* or as *C. rufocanus bedfordiae* (in common name, Bedford's red-backed vole). The grey-sided vole is the most common small mammal in Hokkaido. It inhabits open areas as well as forests, and mainly feeds on green plants. The grey-sided vole has a high reproductive potential; litter size: 4-7; gestation period: 18-19 days; maturation age: 30-60 age in days. Though Spring-born individuals usually attain sexual maturity in their summer-fall of birth, their maturation is sometimes suppressed under high density condition. The breeding season is generally from April to October, but with some regional variation. The vole *C. rufocanus* has a rather specialized diet (folivorous), particularly during winter when it feeds on bamboo grass. Many predators specialize on the grey-sided vole in Hokkaido; even the red fox, which is a typical generalist predator, selectively takes this vole. Damage by voles' barking used to be severe on forest plantations in Hokkaido. Censuses of small rodents have been carried out for management purpose since 1954.

Key words: biogeography, *Clethrionomys rufocanus*, forest damage, life history, monitoring, taxonomy.

Introduction

Cyclic population fluctuations of the grey-sided vole *Clethrionomys rufocanus* (Sundevall, 1846) have fascinated many ecologists, both in Fennoscandia (Kalela 1957, 1971; Viitala 1977), in Siberia (Koshkina 1957, 1965, 1966; Semenov-Tjan-Shanskij 1970) and in Japan (Kinoshita 1928; Ota 1958a, 1968; Ota et al. 1959; Abe 1976a). This species has been intensively studied in Japan both from an applied perspective (forest damage) as well as from the perspective of basic ecology. Ueda et al. (1966) and Ota (1984) provide two comprehensive reviews of the Japanese works. More recent studies on this species include Henttonen et al. (1977, 1987), Saitoh (1981, 1987, 1989, 1990, 1991, 1995), Nakata (1984, 1989), Kawata (1985, 1987, 1988, 1989), Ims (1987a,b,c, 1988, 1989, 1990), Ylönen and Viitala (1987), and Ishibashi et al. (1997). The volume edited by Stenseth (1985a) summarized much of what was known, particularly on the basis of the Western literature, at the time about this species and its congeners.

Besides providing a review of the western literature on the species, the main purpose of this paper is to summarize the Japanese literature on the vole *Clethrionomys rufocanus* and to integrate it with other available studies on the species. Even though some of this literature pertaining to ecological issues have been summarized in English in two recent papers (Bjørnstad et al. 1998a; Saitoh et al. 1998a), we repeat some of this here in order to provide as complete as possible an account of the basic biology of the species and its ecological relationship to its environment. Besides providing a taxonomic account of *C. rufocanus* found in Hokkaido, we describe the initiation and basis for the monitoring program set up over the entire Hokkaido for the purpose of monitoring forest plantations against vole damages. For an exhaustive review of social behavior and social organization, see Ishibashi et al. (1998a,b).

The genus *Clethrionomys*

Roughly speaking, there are, within the family Muridae (Rodentia), three types of body shape: the mouse type which is small with a long tail, the rat type which also has a long tail but is larger than the mouse type, and the vole type which is small with a short tail. The red-backed voles (genus *Clethrionomys*) is of the vole type and belongs to the subfamily Arvicolinae including the genera *Microtus*, *Lemmus* and others (see e.g. Wilson and Reeder 1993). The genus *Clethrionomys* has rooted molar teeth which persist during adult life, being a primitive feature and differing from most other voles with rootless molar teeth, which continue to grow throughout their life. The red-backed vole is, therefore, thought to be less adapted to tough vegetation (Corbet and Harris 1991). Species in this




Fig. 1. Distribution of *Clethrionomys rufocanus* in East Asia. a: Corbet (1978) revised the distribution map of *Clethrionomys rufocanus*, re-identifying specimens that were formerly treated as *Clethrionomys rufocanus* to the species of the genus *Eothenomys*. His examination was, however, insufficient. b: Kaneko (1990, 1992) re-examined the more specimens and revised the distribution map.

genus used to be listed under *Evotomys* (e.g. Hinton 1926) until Palmer (1928) established the priority of *Clethrionomys*. *Clethrionomys* is thought to be most closely related to *Eothenomys* (Iwasa 1998); some authors regard *Clethrionomys* as congeneric with *Eothenomys* (e.g. Corbet 1978; Corbet and Hill 1991).

Clethrionomys contains seven species (Table 1) found widely distributed throughout the northern hemisphere (e.g. Stenseth 1985). These species show great karyotypic similarities (Gamperl 1982; Yoshida et al. 1989; Obara et al. 1995); all have $2n = 56$, with 26 pairs of acrocentric and one pair of metacentric.

Males and females of different species of *Clethrionomys* copulate readily when paired in the laboratory, although they exhibit a preference for conspecific mating when given the choice (Rauschert 1963). Experimental crosses between *C. rutilus* from North America and *C. gapperi* gave no progeny (Matthey 1953; Zimmerman 1965). However, experimental interbreeding of *C. glareolus* and *C. rutilus* from Eurasia (Zimmerman 1965) and of the English *C. glareolus* and the Canadian *C. gapperi* (Grant 1974) produced interspecies hybrids. Female progeny from such crosses were fertile, but male progeny were sterile.

These rodents inhabit cold, mossy, rocky forests and woodlands in both dry and moist areas (Hansson 1985a,b); they also inhabit tundra and bogs. They are active during night and day, summer and winter, scurrying and climbing about stumps, fallen logs and rough-barked trees (e.g. Nowak and Paradiso 1983).

They construct spherical nests of grasses, mosses, lichens, or shredded leaves. These nests are usually hidden under the roots of stumps, logs or brush piles. During winter, globular nests of grass may be placed directly on the ground under the snow, with tunnels radiating from the nest under and in the snow. The diet of *Clethrionomys* consists of tender vegetation, nuts, seeds, bark, lichens, fungi, and insects; they are usually found in forested habitats and are generally omnivorous (Hansson 1985a,b). Food is often stored in the nest for use when the supply is short. When disturbed, red-backed voles may utter a chirplike bark that can be heard from one to two meters away, and they flee or freeze in position, depending on their location and preceding activity.

Taxonomic history

The distribution of *C. rufocanus* is given differently by authors coming from the Far East and Europe (Ognev 1950; Corbet 1978; Kaneko 1990, 1992, 1997). This is due to a very confused situation in the Far East, especially in Korea and North-eastern China (Fig. 1). More specifically this confusion is due to specimens from this region partly has been classified as *Clethrionomys rufocanus* by some workers (Hinton 1926; Ellerman 1941; Ellerman and Morrison-Scott 1951; Ognev 1950; see Kaneko 1990), whereas others have classified them as belonging to the genus *Eothenomys* (Corbet 1978; Kaneko 1990, 1992).

Table 1 : *Clethrionomys* species of the world. Primarily based on Corbet (1978), Hall (1981), Honacki et al. (1982), Gromov and Polyakov (1992), Wilson and Reeds (1993), and Abe et al. (1994) with respect to taxonomy and geographic distribution. Other sources given as footnotes. Modified from the one in Stenseth (1985).

-
- C. californicus* (Merriam, 1890), California red-backed vole. Geographic distribution: Pacific coast coniferous forest from the Columbia River south through W. Oregon & N. California. Population dynamical pattern: unknown. Taxonomical comments: includes *C. mazama* & *C. obscurus*¹; population north of Columbia River are placed in *C. gapperi*.
- C. centralis* (Miller, 1906), Tien Shan forest vole. Geographic distribution: restricted to Tien Shan & Dzhungarsk, Altai Mtns (Russia & Singiang, China). Population dynamical pattern: unknown. Taxonomical comments: referred to in Russia as *C. frater* in current literature. Sometimes *C. centralis* is considered a synonym of *C. glareolus*.
- C. gapperi* (Vigors, 1830), Gapper's red-backed vole. Geographic distribution: Nearctic. From Labrador to N British Columbia (Canada), south in the Appalachians to N Georgia; Great Lakes, Northern Plains, and Rocky Mtns (New Mexico & Arizona) to Columbia River in Washington (USA). Population dynamical pattern: non-cyclic throughout its range^{2,3,4}. Taxonomical comments: includes *C. occidentalis* & *C. caurinus*; closely related to European *C. glareolus* - captive matings produced fertile hybrids but of reduced fertility⁵.
- C. glareolus* (Schreber, 1780); Bank vole. Geographic distribution: Palearctic. W Eurasia, from Britain and Scandinavia south to Pyrenees, Italy, the Balkans, Transcaucasus, and east to central Siberia. Populations on exceptionally high altitude are found in S Norway⁶, at 1400 m a.s.l. and in the French Alps⁷ at 2400 m a.s.l. Population dynamical pattern: both stable and cyclic populations^{4,8}. Taxonomical comments: see *C. gapperi* and *C. centralis*.
- C. rex* Imaizumi, 1971; Darkish red-backed vole. Geographic distribution: restricted to Hokkaido, Japan & other adjunct islets (see Fig. 2). Population dynamical pattern: unknown. Taxonomical comments: includes *C. montanus*. Sometimes *C. rex* is considered a synonym of *C. rufocanus*.
- C. rufocanus* (Sundevall, 1846); Grey-sided vole. Geographic distribution: Palearctic. N Palearctic from Scandinavia to Chukotka and Mongolia, Transbaikalia, NE China; Korea; Kamchatka, Sakhalin (Russia), Hokkaido, Japan & other adjunct islets (see Figs. 1, 2 & 5). Population dynamical pattern: both stable and cyclic populations in Hokkaido^{4,9,10,11}; cyclic in Fennoscandia^{4,12}. Taxonomical comments: includes *C. bedfordiae* and *C. sikotanensis*; one or more separate species may occur in Japan, Korea & China (according to Honacki et al. 1982). Sometimes *C. rex* is considered a synonym of *C. rufocanus*.
- C. rutilus* (Pallas, 1778); Northern red-backed vole. Geographic distribution: Holarctic. N Scandinavia east to Chukotka, south to N Kazakhstan (Russia), Mongolia, Transbaikalia, NE China, Sakhalin and Hokkaido; St. Lawrence Isl. (Bering Sea); Alaska east to Hudson Bay, south to N British Columbia and Manitoba (Canada). Population dynamical pattern: cyclic populations, at least, in Fennoscandia^{3,4,13} and non-cyclic, stable populations in Asia, N America^{3,4,13,14} and W Norway¹⁵. Taxonomical comments: *C. rutilus* and *C. gapperi* are sometimes considered conspecifics.
-

1) Alexander and Verts 1992; 2) Merritt 1981; 3) Henttonen & Hansson 1985a; 4) Henttonen et al. 1985; 5) Grant 1976; 6) Skar et al. 1971; 7) Niethammer & Krapp 1982; 8) Hansson 1979; 9) Bjørnstad et al. 1996; 10) Stenseth et al. 1996; 11) Saitoh et al. 1998b; 12) Koshkina 1966; 13) Henttonen & Hansson 1985b; 14) Whitney 1977; 15) Hansson et al. 1978 (the VøsterÅlen site).

Corbet (1978) first designated the red-backed voles of the Far East as *Eothenomys shanseius*, *E. inez* and *E. eva* in Northeast China and *E. regulus* in Korea; unfortunately he did not go into sufficient details, neither with respect to the description nor with respect to the published distribution maps. Kaneko (1990, 1992) carefully distinguished *Eothenomys* (without roots even during old age) from *Clethrionomys rufocanus* (with roots during old age), on the basis of detailed teeth examination, and clearly showed the southern demarcation of the distribution for *C. rufocanus* (Kaneko 1990, 1992, 1997; Fig. 1).

What in much of the current ecological literature is referred to the grey-sided vole of Hokkaido, is identified as *Clethrionomys rufocanus* by some taxonomists (Corbet 1978; Corbet and Hill 1991;

Musser and Carleton 1993; Abe et al. 1994) or as *C. rufocanus bedfordiae* (in common name, the Bedford's red-backed vole) by other taxonomists (Tokuda 1941; Imaizumi 1960; Abe 1984). It was originally named as a distinct species, *Evotomys* (= now *Clethrionomys*) *bedfordiae* by Thomas (1905a), who mentioned that it is allied to *Evotomys rufocanus* Sundevall, 1846 from Scandinavia, but that *bedfordiae* has a longer tail. Since the roots of the molars develop late in life in *Evotomys rufocanus*, Miller (1900) described the subgenus *Craseomys*. Referring to Miller (1900), Thomas (1905b) changed the name to *Evotomys (Craseomys) bedfordiae* for those specimens having less contrasted general color of red back and grey side, considerably longer tail, and broader and more powerful teeth than *Evotomys rufocanus*. In this paper we will, in order to avoid




Fig. 2. Distribution of *Clethrionomys rufocanus* and *C. rex* in the Hokkaido island and adjunct islets. Shaded areas represent the distribution of *C. rufocanus*. Dots indicate sampling places from which specimens are identified as *C. rex* from 1971 to 1996 (Nakata unpublished). The form of *C. rex* has been reported from Shikotan Is. and Shibotsu Is. shown by asterisks (Kostenko and Arrenoba 1978), though its taxonomic status should be re-examined.

confusion, refer to this vole of Hokkaido as the Bedford's red-backed vole, whereas *C. rufocanus* of the continent is referred to as the grey-sided vole, although we recommend to constantly use the Latin name.

Hinton (1926) arranged the grey-sided vole (then described as one *Evotomys*-species) as *rufocanus*, including *latastei*, *irkutensis*, and *bargusinensis* from Siberia, *shanseius*, *inez*, *nux*, *eva*, *aquilus*, and *alcinous* from Northeast China, *regulus* from Korea, and *smithii*, *rufocanus*, *andersoni*, and *niigatae* from Japan. Then, he revised the vole from Sakhalin, Hokkaido, Honshu, Kyushu and Shikoku as *C. rufocanus smithii* due to the senior synonym. Howell (1929), Allen (1940), Ellerman (1941), Ellerman and Morrison-Scott (1951), Ognev (1950), Jones and Johnson (1965) and Zimmerman (1964) followed this classification.

Kuroda (1931) disagreed with the classification of Hinton (1926) and classified the Bedford's red-backed vole of Hokkaido as *Evotomys* (= now *Clethrionomys*) *rufocanus bedfordiae*, since the Bedford's red-backed vole differs from forms described as *smithii* (*sensu stricto*), *andersoni* (*sensu*

stricto), and *niigatae* (*sensu stricto*) in Honshu, Kyushu, and Shikoku on molar patterns and in the lengths of tail and molar. Since then Tokuda (1941), other Japanese taxonomists and Jameson (1961) followed Kuroda (1931); however, Imaizumi (1972), who did not do so, claimed the Bedford's red-backed vole of Hokkaido to be a distinct species of *C. bedfordiae* having a longer tail, greater skull, smaller auditory bullae, and less conspicuous contrasted color of skin.

Another taxonomic problem of the Bedford's red-backed vole, is that some different forms have been described on small islets off the coast of Hokkaido and alpine areas of Hokkaido. Tokuda (1935) described the new genus and species as *Neoschizomys sikotanensis* from Shikotan, the southern part of Kurile Islands, as a larger form of the red-backed vole (Fig. 2); later he named this vole as *Clethrionomys sikotanensis* (Tokuda 1941). Imaizumi (1949) newly named *Neoschizomys sikotanensis akkessi* from Daikoku Islet off Akkeshi, south-eastern Hokkaido; later he changed the scientific name of the vole from Daikoku and Rishiri Islets to *Clethrionomys sikotanensis* (Imaizumi 1960).




Fig. 3. Upper molar teeth patterns of *Clethrionomys rufocanus* and *C. rex* in a sympatric area of Rishiri Islet. A fine dotted field on the second lingual side on the third upper molar is larger than a black one in *rufocanus*, whereas the size of corresponding areas is similar in *rex* (modified from Kaneko and Sato 1993). CBL: condylobasal length; ML: upper molars length.

Ota (1956) disagreed with the classification of Imaizumi (1949) and demonstrated that one single and distinct species of *C. rufocanus bedfordiae* occurs in Hokkaido and on the Islets of Daikoku and Rishiri. Since rodents are likely to be larger on islands (see Forster 1964), Miyao (1968) followed Ota (1956).

On Rishiri Islet, Imaizumi (1971) described a new form as *Clethrionomys rex* being different from *C. sikotanensis* (larger size, longer tail length and complex molar pattern on the third upper molar). Furthermore, Imaizumi (1972) described a new form of *Clethrionomys montanus* from the Hidaka Mountains, having a longer tail, complex molar pattern on the third upper molar with confluent dental spaces. Corbet (1978) listed *C. rex* as a distinct species from *C. rufocanus*, and considered *montanus* as a conspecific with *rex*. Aimi (1980), however, regarded *C. rex* as synonym with *C. rufocanus*, since the three diagnostic characters described by Imaizumi (1971, 1972) are not found in all specimens on Rishiri Islet. Abe (1984) objected to Aimi (1980) because of too small number of specimens and

characters examined; Abe (1984) concluded, however, that *C. montanus* is conspecific with *C. rex*.

Kaneko and Sato (1993) collected 57 specimens in sympatric distribution of *C. rufocanus* and *C. rex* in a small area of Rishiri Isl. *Clethrionomys rex* has a complex molar pattern on the third upper molar, where the width of salient triangles shows almost the same on both sides. *Clethrionomys rufocanus*, on the contrary, has a large second lingual salient angle shaped round on the third upper molar, where the width of the salient triangle shows larger on the lingual side (Fig. 3). As the increase of CBL (condylobasal length), the molar pattern gradually becomes simple due to smaller posterior loop and appears confluent dental isthmus as found in the holotype of *C. montanus*. They conclude that *Clethrionomys rex* is a distinct species being conspecific with *montanus*. A molecular study also identifies three distinct species of *Clethrionomys* in Hokkaido, namely *C. rufocanus*, *C. rex* and *C. rutilus* (Fig. 4; Wakana et al. 1996).

Judging from the above morphological and molecular findings, we hereafter regard *C. rex* (the darkish red-backed vole) as a distinct species differing from *C. rufocanus* (the grey-sided vole or Bedford's red-backed vole).

Because the holotype of *Neoschizomys sikotanensis* Tokuda, 1935 is lost and the picture of the holotype shows the simple form on the third upper molar (Tokuda 1935), the name of *sikotanensis* is considered to be a junior synonym of *rufocanus*. Therefore, the vole having the complex form on the third upper molar on Shikotan Island seems not to be *C. sikotanensis* (Kostenko and Arrenoba 1978) but *C. rex*. Further examination is needed for the specimens of Shikotan and Shibotsu, the southern part of Kurile Islands, and Sakhalin, where *C. sikotanensis* is distributed (Kostenko and Arrenoba 1978).

Scientific names of genera, species and subspecies related to *Clethrionomys rufocanus* are listed in historical order in Appendix.

Morphology

Clethrionomys rufocanus (Sundevall, 1846)

The vole *Clethrionomys rufocanus* found in Hokkaido is a small-sized, and short-tailed (around 40%) rodent. Data on external dimensions are as follows: head and body length = 107-126 mm, tail length = 44-57 mm, hind foot length = 18.2-20.8 mm, and body weight = 27-50 g (Abe et al. 1994). The sizes varies in Hokkaido and on islets off the coast of Hokkaido (Kaneko et al. unpublished). The fur is dense, long and soft in winter, but shorter and rougher in summer. The general coloration above is dark lead grey with a pronounced reddish wash. The wash becomes less prominent on the sides, which is




Fig. 4. Phylogenetic trees for the mitochondrial DNA haplotypes and major ribosomal DNA repetypes (after Wakana et al. 1996). *Clethrionomys rufocanus* includes *C. sikotanensis*. *Clethrionomys rex* includes *C. montanus*. See the original paper for details.

beige. The underparts are dark slate grey to ivory white. The tail has a slight terminal pencil of hairs. The *C. rufocanus* of Hokkaido has a slightly longer tail than that in Far East, but is almost the same in the hind foot length (Kaneko et al. unpublished).

The skull of adults is massive and angular. Interorbital region is moderately constricted, and parallel-sided. Braincase is large and rectangular. Pterygoid fossa is deep. Auditory bullae is moderately large. The size of adult skulls is larger in *C. rufocanus* from Hokkaido than in *C. rufocanus* from Finland. Data on skull dimensions are as follows: condylobasal length = 26.0-29.8 mm, zygomatic width = 15.0-17.1 mm, nasal length = 7.4-8.7 mm, interorbital width = 3.4-4.3 mm, palatine length = 12.3-13.5 mm, diastema = 7.5-9.2 mm, incisive foramen length = 5.2-6.2 mm, upper molars length = 6.5-7.5 mm, and the length of mandible = 16.8-18.9 mm (Abe 1984). The size varies between the main island of Hokkaido and adjunct islets (Kaneko et al. unpublished).

The size of I-M3 (the distance from the anterior point of incisor to the posterior edge of the third upper molar), where roots of the molars develop, is larger in the vole of Hokkaido (including adjunctive islets) than in voles from the Eurasian continent (e.g. Finland, Mongolia, Northeast China, Northern Korea, and Kamchatka), though the variation in size of I-M3 is overlapping in voles from these regions (Kaneko et al. unpublished). The separate roots developed

slightly earlier in Finland than in Hokkaido: 195 days in Finland (Viitala 1971) versus 200 days in Hokkaido (Abe 1976b). The interorbital width is almost the same in the vole of Hokkaido and Far East (Kaneko et al. unpublished).

Clethrionomys rex and *C. rufocanus*

The darkish red-backed vole *Clethrionomys rex* is dark yellow brown or dark brown on the dorsal side, differing from the red brown on the back of *C. rufocanus* from Hokkaido. External dimensions of *C. rex* are larger than *C. rufocanus* as follows: head and body length = 112-143 mm, tail length = 44-60 mm, hind foot length = 20-22.4 mm and body weight = 33-62 g (Abe et al. 1994). Data on skull dimensions are as follows: condylobasal length = 26.5-29.7 mm, zygomatic width = 15.0-17.4 mm, nasal length = 8.1-9.0, interorbital width = 3.8-4.5 mm, palatine length = 12.8-14.4 mm, incisive foramen length = 5.4-6.4 mm, upper molars length = 6.7-7.6 mm, and the length of mandible - 17.0-19.4 mm (Abe 1984). The complicated pattern on the third upper molar is changed with the increase of CBL and the dentine spaces gradually develops (Kaneko and Sato 1993). The roots of molars develop slower in *C. rex* than in *C. rufocanus* (Abe et al. 1971; Abe 1973); molar roots of *C. rex* are not completely formed even in 250 age in days, whereas they are formed until 200 age in days in *C. rufocanus* (Onoyama et al. unpublished; see also Abe 1973).




Fig. 5. Distribution of *Clethrionomys rufocanus* in the Holarctic (modified from Stenseth 1985). Shaded areas are the distribution formerly suggested, which is revised by Kaneko (1990, 1992). He confirmed the presence of *C. rufocanus* at the places indicated by dots.

Cytology

Even though various *Clethrionomys* species show great karyotypic similarities (Gamperl 1982; Yoshida et al. 1989; Obara et al. 1995), there are at least two karyotypes. A reciprocal translocation between Nos. 1 and 9 chromosomes was first found using G-banding between *C. rufocanus* and *C. glareolus* (Gamperl 1982). Modi and Gamperl (1989) found the same translocation pattern in *C. rutilus*, *C. gapperi* and *C. californicus* as one in *C. glareolus*, and regarded the karyotype of *C. rufocanus* as primitive. This is consistent with the results obtained from traditional morphological analyses (Miller 1900), hybridity analyses (Grant 1974; Zimmermann 1965) and biochemical and molecular genetic analyses (Catzeflis et al. 1987; Tegelström 1987). Although Yoshida et al. (1989) regarded the karyotypes of all species (or forms) of *Clethrionomys* in Hokkaido (*C. rufocanus*, *C. rex*, *C. montanus*, *C. sikotanensis*, and *C. rutilus*) as identical (i.e. *rufocanus*-type), the karyotype of *C. rutilus* in Hokkaido is suggested allocated into be *glareolus*-type by Obara et al (1995). Thus, karyotypes of species (or forms) of *Clethrionomys* in Hokkaido are also classified to two type; *rufocanus*-type: *C. rufocanus*, *C. rex*, *C. montanus*, and *C. sikotanensis*; *glareolus*-type: *C. rutilus* (see also Kashiwabara and Onoyama 1988). According to Wakana et al. (1996), there is no difference in mtDNA between *C. rex* from Rishiri and *C. rex* from Teshio, the latter of which is named as *C. montanus* by the classification of Imaizumi (1972). The electrophoretic patterns of 14 different enzymes and general protein indicate that *C. rufocanus* from Hokkaido is conspecific with that

from Rishiri, which Imaizumi (1971) named *C. sikotanensis* (Yoshida et al. 1989).

The chromosome number of *C. rufocanus* is $2N = 56$ and the number of chromosome arms (FN) is 58. Autosomes of 26 pairs are acrocentric and one pair is metacentric. The X-chromosome is an acrocentric both in continent and in Hokkaido, but Y-chromosome is a submetacentric in Hokkaido but an acrocentric in continent (Vorontsov et al. 1978). The G-band patterns of *C. rufocanus* from Sweden are identical with those of Hokkaido (Gamperl 1982; Mascarello et al. 1974).

The karyotype of *Clethrionomys rex* is $2N = 56$ and $FN = 58$. The autosomes are 26 pairs of acrocentrics and one pair of metacentric. X- and Y-chromosomes are acrocentrics. The G-band patterns are similar between *C. rufocanus* from Hokkaido and *C. rex* from the Hidaka Mountains which is named *C. montanus* according to the classification of Imaizumi (1972) (Kashiwabara and Onoyama 1988).

Distribution of *Clethrionomys rufocanus*

Clethrionomys rufocanus is distributed from Fennoscandia through Siberia (Fig. 5; where the northern Mongolia, Northeast China and Kaima Plateau in North Korea are the southern boundary) to Kamchatka, Sakhalin and Hokkaido (Ognev 1950; Henttonen and Viitala 1982; Kaneko 1990, 1992, 1997). *Clethrionomys rufocanus bedfordiae* is distributed in Hokkaido, on islets off Hokkaido such as Rishiri, Rebun, Daikoku, Teuri, and Yagishiri, and the southern part of Kurile Islands (i.e. Kunashiri,




Fig. 6. Frequency distributions of litter size. a.: Frequency distribution of the number of embryos obtained from 747 pregnant females sampled at various sites in Hokkaido between 1967 and 1974 (Fujimaki 1981); average = 5.27. b.: Frequency distribution of the number of litter mates obtained from 123 litters in an experimental population in Sapporo, Hokkaido between 1992 and 1993 (Saitoh and Ishibashi unpublished data); average = 4.79. Litter mates were determined based on the following information: DNA analyses for parentage, reproduction career of a mother candidate, and date and site at the first capture of a juvenile.

Shikotan, Shibotsu and others; Fig. 2; Ota 1956; Kostenko and Arrenoba 1978; Kaneko et al. unpublished).

Clethrionomys rex is distributed in Hokkaido (Daisetsu, Hidaka, Teshio, Kitami and Yuubari Mountains, Abuta Town near Lake Tooya; Fig. 2; Nakata unpublished), islets off Hokkaido such as Islets of Rebun and Rishiri (Abe et al. 1994), Shikotan and Shibotsu, the southern part of Kurile Islands, and Sakhalin (Kostenko and Arrenoba 1978).

The ecology of *Clethrionomys rufocanus* of Hokkaido

Growth and reproduction

The gestation period of the Bedford's red-backed vole is 18-19 days under benign conditions (Abe 1968). The litter size is typically 4-7 (Fig. 6; Kinoshita 1928; Fujimaki 1972, 1973, 1975, 1981; Kuwahata 1984; Nakata 1984; Saitoh and Ishibashi unpublished; see Table 1 in Ueda et al. 1966 for review). Infants are born weighing about 2.0 g and grow fast; they wean after about 17 days and reach 22-23 g around an age of 30 days. Age of maturation for *C. rufocanus* is around 30-60 days under benign conditions (Abe 1968), but growth pattern varies between different generations: spring-born individuals grow faster and become mature earlier than autumn born ones (Kinoshita 1928; Ota et al. 1961). Spring-born individuals attain sexual maturity in the year they are born. Autumn-born voles remain reproductively inactive through the winter (Fujimaki 1975; Kuwahata 1984). The maturation of females is influenced by social interaction in a natural condition (Saitoh 1981, 1990). Spring populations are composed of overwintered animals. The proportion of this cohort declines to 10% or less by October/November (Fujimaki 1969a, b, 1975; Abe 1976a, b; Fujimaki et al. 1976; Kuwahata 1984). Generally, the lifetime of free ranging individuals is less than one year (Yoneda 1982).

The breeding season

The breeding season is generally from April to October. Although pregnant females are common between May and October, pregnancy rate is higher in spring and autumn than in summer; the proportion of mature males also is low during the middle of the summer (Ota et al 1959; Kinoshita and Maeda 1961; Kuwahata 1962). Thus, the breeding activity has two peaks (spring and autumn) in a year. Most breeding females give birth to 2-4 litters through the breeding season (spring or autumn) and post partum estrous is frequently observed in spring (Saitoh and Ishibashi unpublished data). Spring-born individuals typically mature during the summer and comprise a major part of the breeding autumn population. Late summer- or autumn-born individuals generally wait until the next spring.

In contrast to this general breeding pattern, populations in high elevation or in the eastern area, where temperature (particularly that during the summer) is lower than in the lowland are reported to have higher pregnancy rates during the summer than that during the autumn (Ueda 1961). This leads to a single peak in breeding activity through the season, rather than the two peaks described above. It appears, thus, that temperature influences the breeding activity. A single peak in the breeding pattern is also reported during periods of high densities in

populations near Sapporo, where the two-peak pattern is the norm (Fujimaki 1969; Abe 1976b).

There is also some other geographic changes in the breeding pattern. In the southern part of Hokkaido where temperature is higher and the snow disappears earlier, the breeding starts in the middle of March, and early spring-born individuals may become mature in June. In the colder northern and eastern parts of Hokkaido, the breeding starts in April (Fujimaki 1972). However, since voles in the northern or eastern areas become mature at a smaller size, autumn breeding still tends to be more active in these areas (Fujimaki 1973).

Even though winter breeding is known to occur (Kaikusalo and Tast 1984; Nakata 1987; Saitoh 1989), there is no evidence to expect it to be quantitatively important in Hokkaido.

Factors influencing breeding

Population density: High population density suppresses the maturation of young females (Saitoh 1981). Ota and Fujikura (1959) and Kuwahata (1966) reported low maturation rate of young individuals in high density years. The breeding pattern with one peak a year near Sapporo (see above) may be caused by high populating densities, and not by temperature (Abe 1976a). The breeding season is shorten in a high-density year but elongated in a low-density year (Nakata 1989). High densities may also lead to reduction of litter size (Nakata 1984).

Fluctuation phase: Even though under similar densities, reproductive activity varies between different fluctuation phases. The proportion of breeding voles is higher during the increase phases than during the decline phases (Nakata 1989). Litter size also varies among phases with a similar pattern (Nakata 1984).

Food conditions: The breeding is usually more active in forests plantations than in natural forests, because the undergrowth in plantations is generally rich (Kinoshita and Maeda 1961; Maeda 1963). This is particularly so in young plantations, in which green plants are growing well owing to more sunlight as compared to those in mature natural forests. If provided ad libitum food, voles will breed in outdoor enclosure or cages in the winter (Haga 1954; Maeda 1962; Saitoh 1989). Thus, there is evidences that quality and quantity of food greatly influence the breeding activity (see also Andreassen and Ims 1990).

Habitat and diet

The Bedford's red-backed vole is a habitat generalist

in Hokkaido. Although the species of the genus *Clethrionomys* generally favor forests and woodlands, the vole inhabits open habitats as well as forests in Hokkaido, possibly because there are no *Microtus* (Ota and Jameson 1961). The vole is the most strictly ground-dwelling among rodents in Hokkaido (Ota 1968; Abe 1986; Abe et al. 1989). That is, it is much more ground-dwelling than the three sympatric species of *Apodemus*. A dominant understory plant species in forests is the bamboo grass (*Sasa* spp.; Wada 1993). Vegetation cover and density of understory vegetation are well developed from mid June to October (Yoneda 1983). The snow rich period is associated with relatively high survival in *C. rufocanus* (Dewa 1975 see also Yoccoz et al. 1998). There is also some evidence (Yoneda 1983), that dense vegetation conveys some of the same protection against hostility factors such as predators or unfavorable weather conditions.

The Bedford's red-backed vole mainly feeds on green plants, which usually occupy more than 50% of its diet (Ota et al. 1959, Ota 1984). Leaves and shoots of bamboo grass dominate the food during winter, while various forbs and grasses are eaten during summer (Ota 1984). The diet is, thus, predominantly folivorous and subdominantly granivorous. The grey-sided vole is more folivorous than the other *Clethrionomys* species (cf. Hansson 1985). This feeding habit is particularly prevalent in Hokkaido - possibly due to the absence of *Microtus*. However, acorns are also, to some extent, eaten in the autumn while bark is taken during winter (Ota 1984).

Competitors and predators

Species of *Microtus* seem to be superior competitors for the grey-sided vole (*C. rufocanus*) in the Eurasian continent (Henttonen et al. 1977; Hanski and Henttonen 1996). However, there are no *Microtus* species in Hokkaido. Although two species of the wood mouse (*Apodemus argenteus* and *A. speciosus*) are common and sympatric with the Bedford's red-backed vole, there is no evidence of competition between *Clethrionomys* and *Apodemus* (Abe 1986). Congeneric interactions are, however, thought to occur (Ota 1968; Abe 1986; Nakata 1995). A possible competitor, the northern red-backed vole (*C. rutilus*) is inferior to the Bedford's red-backed vole in most habitats, except for in pure coniferous forests (Ota 1984). Habitat segregation is observed between *C. rufocanus* and *C. rex* (Kaneko and Sato 1993; Nakata 1995).

Three mustelid species are important predators for small rodents in Hokkaido (in intuitive order of importance): *Mustela nivalis*, *M. itatsi*, and *M. vison*. *Martes zibellina* and *Mustela erminea* are certainly also present but there is little information about these species. The red fox (*Vulpes vulpes*), two owls (*Strix uralensis* and *Asio otus*) and three species of snakes (*Elaphe climacophora*, *E. conspicillata*, *E.*




Fig. 7. Yearly changes in areas of forestry plantations and the damaged areas of the plantations by the vole in hectares.

quadrivirgata and *Agkistrodon blomhoffii*) are also important predators for rodents (Kadowaki personal communication). Predators prefer the Bedford's red-backed vole to the *Apodemus* species in Hokkaido; even the red fox, which is a typical generalist predator, selectively takes this vole (Yoneda 1979). The reported percentages of occurrence in dropping or stomach contents of the two genera are: 30.7% vs 0.3% in *M. vison* (Uraguchi et al. 1987), 30-60% vs 1% in *V. vulpes* (Abe 1975, Misawa 1979, Kondo et al. 1986), 25%-50% vs 5%-25% in *S. uralensis* (Matsuoka 1977, Yoneda et al. 1979) and 98-100% vs 0.8-3.7% in *A. otus* (Matsuoka 1974).

Although many parasites are known in *C. rufocanus* of Hokkaido, its influence on population dynamics is uncertain (Yoneda and Nakata 1984; see also Saitoh and Takahashi 1998).

Forest management and rodent damage

Hokkaido is a forest-rich island: forests once covered about 70% of area of the island. Forestry plantations were started in 1886 and forest damage by vole-barking was first reported in 1904 in Hokkaido (*Rodent Research Group* 1958). Scientific investigations of both applied and basic questions pertaining to the ecology of the Bedford's red-backed vole in Hokkaido were initiated by Kinoshita (1928, see also Ueda et al. 1966). Thereafter, because of severe damage spreading all over Hokkaido (e.g. in 1937, 1942-43, and 1951), such studies were encouraged and have been followed up by researchers in *Hokkaido University*, *Forestry and Forest Products Research Institute*, and *Hokkaido*

Forestry Research Institute.

The Forestry Agency of Japanese Government (FAJ) instigated forestry plantations since 1950s to transform natural forests into more productive plantations. Although forest planting followed various silvicultural procedures, the most common procedure, which was adopted in Hokkaido, was the drastic method of transforming a natural system into an artificial one. After clearance of natural forests, weeds were removed from the clear-cut, then larch *Larix kaempferi* or todo fir *Abies sachalinensis* seedlings were planted densely. Because weeds grow thickly in young plantations and may suppress the growth of tree seedlings, the weeds are cut every summer for the next five to nine years after tree planting. This procedure have provided a favorable habitat for the Bedford's red-backed vole (Ota 1968; Saitoh and Nakatsu 1997), just like clearcutting appear to favor folivorous rodents elsewhere (Hansson 1992).

This forestry practice was implemented faithfully, all over Hokkaido. Considerable areas of natural forests were clear-cut and transformed into single species, largely coniferous, plantations (Fig. 7). During the peak of this development, more than one percent of natural forests were cut and transformed into coniferous plantations every year. Because this intensive planting extended the damage by the Bedford's red-backed vole, FAJ in Hokkaido and Hokkaido Government initiated a management-focused census program of small rodents, together with *The Rodent Research Group*, which was organized in 1953 by researchers of in Hokkaido University, *Forestry and Forest Products Research*




Fig. 8. Distribution of forests in Hokkaido and of census locations in national forests. Dense shaded area: national forests managed by the Forestry Agency of Japanese Government (FAJ); slightly dense shaded areas: prefectural forest managed by Hokkaido Government; light shaded areas: private forests; solid areas: experimental forests of universities. Circles indicate locations of Ranger Offices in national forests, which carried out censuses of rodents.

Institute, and Hokkaido Forestry Research Institute. After the preliminary work in the early 1950s aiming at devising an appropriate census method, census methods using snap traps were standardized in 1954. The methods were standardized again into using 50 snap traps at 10-m grid pattern in 1956, which have been used until the present (Fujimaki 1977; for details of methods, Saitoh et al. 1997, 1998a,b).

There are two census programs implemented by FAJ in Hokkaido and Hokkaido Government using the same trapping protocol; trapping three times a year (spring, summer and fall) in selected habitats in forests all over Hokkaido by foresters; both in young plantations (until about 15 years after planting) and in natural forests neighboring the plantations (Fig. 8). In the case of national forests managed by FAJ, Ranger Offices (which usually manage several thousands hectares of forests) were the unit in the census; e.g. a total of 436 Ranger Offices (which have at least two census sites in a plantation and in a natural forest) carried out censuses in 1992. The personnel of Ranger Offices, who were regularly trained by specialists, trapped and identified rodents. These are the data used for the spatio-temporal analysis reported by Saitoh et al. (1998) and by Bjørnstad et al. (1998) later in this special issue.

In private and prefecture's forests the Hokkaido Government has carried out three types of censuses. For private forests, there are furthermore two types; one is the census by designated trained specialists whereas another is carried out by foresters. In prefecture's forests trained personnel of these offices perform the censuses. Most private forests are near suburban or cultivated areas ranging from lowland to mountainous areas, whereas most national and prefecture's forests are generally located in mountainous areas (Fig. 8).

Ota (1958b) first predicted, on the basis of the census data, a vole outbreak for the following year and the *Rodent Research Group* gave foresters a warning of damage. However, they were not able to check the unprecedented damage in 1959, when more than 60 million saplings were destroyed (Fig. 7). Thereafter, using the census data systematically, density in autumn came to be estimated from density in spring or summer. Based on these estimates and the past record on damages, the amount of poisonous baits to be scattered through forestry plantations was determined (Nakata 1986).

That census data has also proven invaluable for population ecological studies (Krebs 1997). Unfortunately, the census efforts were greatly

reduced during the middle of 1990s; this coincided with a reduction of the area with new forestry plantations - a development, which began during late 1980s. This is caused mainly by the lack of appropriate areas for such plantations, by economic hardship of the forestry organizations and by public demands for less clearcutting; forestry plantations are usually made on areas where natural forests were clear-cut. However, data till 1992 have provided us with a most valuable set of data for describing and analyzing population fluctuations of the Bedford's red-backed vole. Parts of the census data, covering the northern national forests, have already been analyzed in some detail (Saitoh 1987; Bjørnstad et al. 1996, 1998a; Stenseth et al. 1996; Saitoh et al. 1997, 1998a,b). Saitoh et al. (1998b; see also Bjørnstad et al. 1998b) summarizes and extends these analyses to the entire set of time series from the census program in the national forests surveyed by FAJ. This set of time series in Hokkaido certainly represents a goldmine for spatio-temporal analyses (Krebs 1997). Fluctuation patterns found using these data, together with the rich background information on the biology of the Bedford's red-backed vole reviewed in this Special Feature papers, provide a unique opportunity for understanding population fluctuations in rodents. Further work on the analysis of the monitoring data should be coupled with more detailed (the challenging) work on the demography and social organization of the Bedford's red-backed voles populations in the various regions of Hokkaido.

Acknowledgements: We are indebted to Hisashi Abe, Masahiro Iwasa, Heikki Henttonen, Per Erik Jorde and Kazimierz Kowalski for having provided information to and commented on earlier versions of this paper.

References

- Abe, H. (1968) Growth and development in two forms of *Clethrionomys*. 1. External characters, body weight, sexual maturity and behavior. *Bulletin of the Hokkaido Forest Experiment Station* (Bibai) **6**: 69-89 (in Japanese with English summary).
- Abe, H. (1973) Growth and development in two forms of *Clethrionomys*. II. Tooth characters, with special reference to phylogenetic relationships. *Memoirs of the Faculty of Agriculture, Hokkaido University* **57**: 229-254.
- Abe, H. (1975) Winter food of the red fox, *Vulpes vulpes schrencki* Kishida (Carnivora: Canidae), in Hokkaido, with special reference to vole populations. *Applied Entomology and Zoology* **10**: 40-51.
- Abe, H. (1976a) Population structure and reproductive activity of *Clethrionomys rufocanus bedfordiae* (Thomas) in a wind shelter-belt of the Ishikari plain, Hokkaido. *Journal of the Mammalogical Society of Japan* **7**: 17-30 (in Japanese with English summary).
- Abe, H. (1976b) Age determination of *Clethrionomys rufocanus bedfordiae* (Thomas). *Japanese Journal of Ecology* **26**: 221-227 (in Japanese with English summary).
- Abe, H. (1984) Taxonomy on wild murid rodents in Hokkaido. pp. 1-20. In Ota, K.(ed.) *Study on wild murid rodents in Hokkaido*. Hokkaido University Press, Sapporo. (in Japanese)
- Abe, H. (1986) Vertical space use of voles and mice in woods of Hokkaido, Japan. *Journal of the Mammalogical Society of Japan* **11**: 93-106.
- Abe, H., T. Kobayashi, K. Maeda and I. Hayata (1971) Results of the small mammal survey on the Daisetsu Area. *Annual Report JIBP/CT-S Fisc. 1970*:13-22.
- Abe, H., T. Shida, and T. Saitoh (1989) Effects of reduced vertical space and arboreal food supply on densities of three rodent species. *Journal of the Mammalogical Society of Japan* **14**: 43-52.
- Abe, H., N. Ishii, Y. Kaneko, K. Maeda, S. Miura and M. Yoneda (1994) *A pictorial guide to the mammals of Japan*. Tokai University Press, Tokyo. (in Japanese)
- Aimi, M. (1980) A revised classification of the Japanese red-backed voles. *Memoirs Faculty of Science, Kyoto University, Series of Biology* **8**: 35-84.
- Alexander, L. F. and B. J. Verts (1992) *Clethrionomys californicus*. *Mammalian Species* **406**: 1-6.
- Allen, G. M. (1940) Mammals of China and Mongolia. Pt. 2. Publ. Amer. Mus. Nat. Hist., Central Asiatic Exped., **11**: 621-1350.
- Andreassen, H. P. and R. A. Ims (1990) Responses of female grey-sided voles *Clethrionomys rufocanus* to malnutrition: a combined laboratory and field experiment. *Oikos* **59**:107-14.
- Bjørnstad, O. N., S. Champely, N. C. Stenseth and T. Saitoh (1996) Cyclicity and stability of grey-sided voles, *Clethrionomys rufocanus*, of Hokkaido: spectral and principal components analyses. *Philosophical Transactions of the Royal Society of London B* **351**: 867-875.
- Bjørnstad, O. N., N. C. Stenseth and T. Saitoh (1998a) Synchrony and scaling in dynamics of voles and mice in northern Japan. *Ecology* (in press)
- Bjørnstad, O. N., N. C. Stenseth, T. Saitoh and O. C. Lingjære (1998) Mapping the regional transitions to cyclicity in *Clethrionomys rufocanus*: spectral densities and functional data analysis. *Researches on Population Ecology* **40**: 77-84.
- Corbet, G. B. (1978) *The mammals of the Palaearctic region: a taxonomic review*. Cornell University Press, London-Ithaca.

- Corbet, G. B. and J. E. Hill (1991) *A world list of mammalian species*. 3rd. edn. British Museum (Natural History), London.
- Corbet, G. B. and S. Harris (1991) *The handbook of British mammals*. 3rd. edn. Blackwell Scientific Publications, London.
- Dewa, H. (1975) Seasonal variation of daily activity rhythms in snow season. *Research Bulletin of College Experimental Forests Hokkaido University (Sapporo)* **22**: 105-120. (in Japanese with English summary)
- Ellerman, J. R. (1941) *The families and genera of living rodents. Vol. 2. Family Muridae*. British Museum (Natural History), London.
- Ellerman, J. R. and T. C. S. Morrison-Scott (1951) *Checklist of palaeartic and Indian mammals, 1758 to 1946*. British Museum (Natural History), London.
- Forster, J. B. (1964) Evolution of mammals on islands. *Nature* **202**: 234-235.
- Fujimaki, Y. (1969a) Reproductive activity in *Apodemus argenteus* Temminck. *Journal of the Mammalogical Society of Japan* **4**: 74-80. (in Japanese with English summary)
- Fujimaki, Y. (1969b) The fluctuations in the number of small rodents. *Bulletin of the Hokkaido Forest Experiment Station* **7**:62-77. (in Japanese with English summary)
- Fujimaki, Y. (1972) Reproductive activity in *Clethrionomys rufocanus bedfordiae* 1. Regional differences in spring reproductive activity. *Bulletin of the Hokkaido Forest Experiment Station* **10**:59-67. (in Japanese with English summary)
- Fujimaki, Y. (1973) Reproductive activity in *Clethrionomys rufocanus bedfordiae* 2. Regional differences in summer reproductive activity. *Bulletin of the Hokkaido Forest Experiment Station* **11**: 122-131. (in Japanese with English summary)
- Fujimaki, Y. (1975) Reproductive activity in *Clethrionomys rufocanus bedfordiae* 3. Regional differences in autumn reproductive activity. *Bulletin of the Hokkaido Forest Experiment Station* **13**:38-45. (in Japanese with English summary)
- Fujimaki, Y. (1977) Historical transition in census methods for small rodents and in prediction systems for vole density. *Nonezumi (Voles and Mice)* **142**: 74-75. (in Japanese)
- Fujimaki, Y. (1981) Reproductive activity in *Clethrionomys rufocanus bedfordiae* 4. Number of embryos and prenatal mortality. *Japanese Journal of Ecology* **31**:247-256.
- Fujimaki, Y., S. Mizushima, and H. Dewa (1976) Age determination in two species of *Apodemus*. *Japanese Journal of Ecology* **26**:19-23 (in Japanese with English summary).
- Gamperl, R. (1982) Chromosomal evolution in the genus *Clethrionomys*. *Genetica* **57**: 193-197.
- Grant, P. R. (1974) Reproductive compatibility of voles from separate continents (Mammalia: *Clethrionomys*). *Journal of Zoology* **174**: 245-254.
- Grant, P. R. (1976) An 11-year study of small mammal populations at Mount St. Hilaire, Quebec. *Canadian Journal of Zoology* **54**: 2156-2173.
- Gromov, I. M. and I. Ya. Polyakov (1992) Voles (Microtinae). Fauna of the USSR Mammals, volume III, No. 8. E. J. Brill, Leiden.
- Haga, R. (1954) The behavior of the vole in the snow season. *Memoirs of the Faculty of Agriculture, Hokkaido University* **2**: 66-78. (in Japanese with English summary)
- Hall, E. R. (1981) *The mammals of North America*. John Wiley and Sons, New York.
- Hansson, L. (1974) Nya utbredningsuppgifter för grÅsiding, med synpunkter på eventuell konkurrens med skogssork. *Fauna och Flora* **69**:81-112. (in Swedish)
- Hansson, L. (1979) Food as a limiting factor for small rodent numbers. *Oecologia* **37**:297-341.
- Hansson, L. (1985a) *Clethrionomys* food: generic, specific and regional characteristics. *Annals Zoologici Fennici* **22**:315-318.
- Hansson, L. (1985b) The food of bank voles, wood mice and yellow-necked mice. *Symposium of the Zoological Society of London* **55**:141-168.
- Hansson, L. (1992) Small mammal communities on clearcuts in a latitudinal gradient. *Acta Oecologia* **13**:687-699
- Hansson, L. and H. Henttonen (1985a) Gradients in density variations of small rodents: the importance of latitude snow cover. *Oecologia* **67**:394-402.
- Hansson, L. and H. Henttonen (1985b) Regional differences in cyclicity and reproduction in *Clethrionomys* species: Are they related? *Annales Zoologica Fennici* **22**: 277-288.
- Hansson, L., J. Lofqvist and A. Nilsson (1978) Population fluctuations in insectivores and small rodents in northernmost Fennoscandia. *Zeitschrift für Säugetierkunde* **43**: 75-92.
- Henttonen, H. and J. Viitala. (1982) *Clethrionomys rufocanus* (Sundevall, 1846)- Graurotelmaus. pp. 147-164. In J. Niethammer and F. Krapp (eds.) *Handbuch der Säugetiere Europas*. Vol. 2. Akademische Verlags, Wiesbaden.
- Henttonen, H., A. Kaikusalo, J. Tast and J. Viitala (1977) Interspecific competition between small rodents in subarctic and boreal ecosystem. *Oikos* **29**:

581-590.

Henttonen, H., A. D. McGuire, and L. Hansson (1985) Comparisons of amplitudes and frequencies (spectral analyses) of density variations in long-term data sets of *Clethrionomys* species. *Annales Zoologici Fennici* **22**:221-227.

Henttonen, H., T. Oksanen, A. Jortikka and V. Haukisalml (1987) How much do weasels shape microtine cycles in the northern Fennoscandia taiga? *Oikos* **50**: 353-365.

Henttonen, H., L. Hansson, and T. Saitoh (1992) Rodent dynamics and community structure: *Clethrionomys rufocanus* in northern Fennoscandia and Hokkaido. *Annales Zoologici Fennici* **29**:1-6.

Honacki, J. H., K. E. Kinman and J. W. Koepf (eds. 1982) *Mammal species of the world. A taxonomic and geographic reference*. Allen Press, Inc., New York.

Howell, A. B. (1929) Mammals from China in the collections of the United States National Museum. *Proceedings of the United States National Museum* **75**: 1-82.

Hinton, M. A. C. (1926) *Monograph of voles and lemmings. Living and extinct*. vol. 1 British Museum (Natural History), London.

Imaizumi, Y. (1949) *The Natural History of Japanese Mammals*. Yoyoshobo, Tokyo. (in Japanese)

Imaizumi, Y. (1960) *Coloured illustrations of the mammals of Japan*. Hoikusha, Osaka. (in Japanese)

Imaizumi, Y. (1971) A new vole of the *Clethrionomys rufocanus* group from Rishiri Island, Japan. *Journal of the Mammalogical Society of Japan* **5**: 99-103.

Imaizumi, Y. (1972) Land mammals of the Hidaka Mountains, Hokkaido, Japan, with special reference to the origin of an endemic species of the genus *Clethrionomys*. *Memoirs of the National Science Museum, Tokyo* **5**:131-149. (in Japanese with English summary)

Ishibashi, Y., T. Saitoh, S. Abe and M. C. Yoshida (1997) Sex-related spatial kin structure in a spring population of grey-sided voles *Clethrionomys rufocanus* as revealed by mitochondrial and microsatellite DNA analyses. *Molecular Ecology* **6**: 63-71.

Ishibashi, Y., T. Saitoh and M. Kawata (1998a) Social organization of the vole *Clethrionomys rufocanus* its demographic and genetic consequences: a review. *Researches on Population Ecology* **40**: 39-50.

Ishibashi, Y., T. Saitoh, S. Abe and M.C. Yoshida (1998b) Kin-related social organization in a winter

population of the vole *Clethrionomys rufocanus*. *Researches on Population Ecology* **40**: 51-59.

Ims, R. A. (1987a) Differential reproductive success in a peak population of the grey-sided vole *Clethrionomys rufocanus*. *Oikos* **50**: 103-113.

Ims, R. A. (1987b) Responses in spatial organization and behaviour to manipulations of the food resource in the vole *Clethrionomys rufocanus*. *Journal of Animal Ecology* **56**: 585-596.

Ims, R. A. (1987c) Determinants of competitive success in *Clethrionomys rufocanus*. *Ecology* **68**: 1812-1818.

Ims, R. A. (1988) Spatial clumping of sexually receptive females induces space sharing among male voles. *Nature* **335**: 541-543.

Ims, R. A. (1989) Kinship and origin effects on dispersal and space sharing in *Clethrionomys rufocanus*. *Ecology* **70**: 607-616.

Ims, R. A. (1990) Determinants of natal dispersal and space use in grey-sided voles, *Clethrionomys rufocanus*: a combined field and laboratory experiment. *Oikos* **57**: 106-113.

Iwasa, M. (1998) Chromosomal variation and molecular phylogeny in red-backed voles. *Honyurui Kagaku* (Mammalian Science) **37**: in press. (in Japanese with English summary)

Jameson, E. W. Jr. (1961) Relationship of the red-backed voles of Japan. *Pacific Science* **15**: 594-604.

Jones, J. K. and Johnson, D. H. (1965) Synopsis of the lagomorphs and rodents. *The University of Kansas Publications of the Museum of Natural History* **16**: 357-407.

Kaikusalo, A. and J. Tast. (1984) Winter breeding of microtine rodents at Kilpisjärvi, Finnish Lapland. *Special Publication of the Carnegie Museum of Natural History* **10**: 243-52.

Kalela, O. (1957) Regulation of reproductive rate in subarctic populations of the vole, *Clethrionomys rufocanus* (Sund.). *Annales Academiae Scientiarum Fennicae, Series A, IV Biologica* **34**: 1-60.

Kalela, O. (1971) Seasonal trends in the sex ratio of the grey-sided vole, *Clethrionomys rufocanus* (Sund.). *Annales Zoologici Fennici* **8**:425-455.

Kaneko, Y. (1990) Identification and some morphological characters of *Clethrionomys rufocanus* and *Eothenomys regulus* from USSR, Northeast China, and Korea in comparison with *C. rufocanus* from Finland. *Journal of Mammalogical Society of Japan* **14**: 129-148.

Kaneko, Y. (1992) Identification and some morphological characters of *Clethrionomys rufocanus*, and *Eothenomys shansei*, *E. inez* and *E. eva* from USSR, Mongolia, and Northeast and central

- China. *Journal of Mammalogical Society of Japan* **16**: 71-95.
- Kaneko, Y. (1997) Classification, geographical distribution and speciation in some rodent species of Murinae and Arvicolinae subfamilies. *Honyurui Kagaku* (Mammalian Science) **37**: 55-74. (in Japanese with English summary)
- Kaneko, Y. and M. Sato (1993) Identification and distribution of red-backed voles from Rishiri Isl., Hokkaido (preliminary study). *Rishiri Town Museum Annual Report* **12**: 37-47. (in Japanese)
- Kawata, M. (1985) Mating system and reproductive success in a spring population of the red-backed vole, *Clethrionomys rufocanus bedfordiae*. *Oikos* **45**: 181-190.
- Kawata, M. (1987) Pregnancy failure and suppression by female-female interaction in enclosed populations of the red-backed vole, *Clethrionomys rufocanus bedfordiae*. *Behavioral Ecology and Sociobiology* **20**: 89-97.
- Kawata, M. (1988) Mating success, spatial organization, and male characteristics in experimental field populations of the red-backed vole, *Clethrionomys rufocanus bedfordiae*. *Journal of Animal Ecology* **57**: 217-235.
- Kawata, M. (1989) Growth and dispersal timing in male red-backed voles *Clethrionomys rufocanus bedfordiae*. *Oikos* **54**: 220-226.
- Kashiwabara, S. and K. Onoyama (1988) Karyotypes and G-banding patterns of the red-backed voles, *Clethrionomys montanus* and *C. rufocanus bedfordiae* (Rodentia, Microtinae). *Journal of the Mammalogical Society of Japan* **13**: 33-41.
- Kinoshita, E. (1928) Study on voles and mice from the standpoint of forest protection. *Research Bulletins of the College Experiment Forests, College of Agriculture, Hokkaido University* (Sapporo) **5**: 1-115. (in Japanese)
- Kinoshita, E. and M. Maeda (1961) Field experiments on the biology of field mice in the woodland. *Bulletin of the Forestry and Forest Products Research Institute* (Ibaraki) **127**: 61-98. (in Japanese with English summary)
- Kondo, N., K. Takahashi, and K. Yagi. 1986. Winter food of the red fox, *Vulpes vulpes schrencki* Kishida, in the endemic area of multilocular echinococcosis. *Bulletin of Preparative Office of Nemuro Municipal Museum* **1**: 23-31 (in Japanese with English summary).
- Koshkina, T. V. (1957) Comparative ecology of the red-backed voles in the northern taiga. *Materials about rodents* **5**: 3-65. (in Russian, edited and translated by W. A. Fuller)
- Koshkina, T. V. (1965) Population density and its importance in regulating the abundance of the red vole. *Bulletin of the Moscow Society of Naturalists, Biological Section* **71**: 5-19. (in Russian)
- Koshkina, T. V. (1966) On the periodical changes in the numbers of voles (as exemplified by the Kola Peninsula). *Bulletin of the Moscow Society of Naturalists, Biological Section* **71**: 14-26. (in Russian)
- Kostenko, B. A. and T. B. Arrenoba. (1978) Characteristics of morphology and biology in the red-backed vole (*Clethrionomys*) on Shikotan Island. *Ecology and Zoogeography of Terrestrial Vertebrates in Far East* 119-125. (in Russian)
- Krebs, C. J. (1997) Vole cycle on Hokkaido: a time-series goldmine. *Trends in Ecology and Evolution* **12**: 340-341.
- Kuroda, N. (1931) A new locality and scientific name of red-backed vole, *Evotomys niigatae*. *Dobutsugaku Zasshi* (Zoological Magazine) **43**: 661-666. (in Japanese)
- Kuwahata, T. (1962) Studies on population fluctuation of the red-backed vole, *Clethrionomys rufocanus bedfordiae* (Thomas) (I). An analysis of a process of the gradation. *Bulletin of the Forestry and Forest Products Research Institute* (Ibaraki) **143**: 15-38. (in Japanese with English summary)
- Kuwahata, T. (1966) Studies on population fluctuation of the red-backed vole, *Clethrionomys rufocanus bedfordiae* (Thomas) (II). Reproductive activity. *Annual Report of the Hokkaido Branch, the Forestry and Forest Products Research Institute* (Sapporo) **143**: 210-236. (in Japanese with English summary)
- Kuwahata, T. (1984) Studies on reproduction and population dynamics of the red-backed vole, *Clethrionomys rufocanus bedfordiae* (Thomas). *Bulletin of the Forestry and Forest Products Research Institute* (Ibaraki) **327**: 1-81. (in Japanese with English summary)
- Maeda, M. (1963) Field experiment on the biology of field mice in woodland 2. Natality and mortality of the red-backed vole, *Clethrionomys rufocanus bedfordiae* (Thomas). *Bulletin of the Forestry and Forest Products Research Institute* (Ibaraki) **160**: 1-18. (in Japanese with English summary)
- Matthey, R. (1953) Les chromosomes des Muridae. *Revue Suisse de Zoologie* **60**: 225-283.
- Matsuoka, S. (1974) Prey taken by long-eared owl *Asio otus* in the breeding season in Hokkaido. *Miscellaneous Reports of the Yamashina Institute of Ornithology* **41**: 324-329. (in Japanese with English summary)
- Matsuoka, S. (1977) Winter food habits of the Ural owl *Strix uralensis* Pallas in the Tomakomai

- experiment forest of Hokkaido University. *Research Bulletins of the College Experiment Forests, Faculty of Agriculture, Hokkaido University* (Sapporo) **34**:161-174. (in Japanese with English summary)
- Mascarello, J. T., A. D. Stock and S. Pathak (1974) Conservatism in the arrangement of genetic material in rodents. *Journal of Mammalogy* 695-704.
- Miller, G. S., Jr. (1900) Preliminary revision of the European redbacked mice. *Proceedings of the Washington Academy of Sciences* **2**: 83-109.
- Merritt, J. F. (1981) *Clethrionomys gapperi*. *Mammalian Species* 146: 1-9.
- Miyao, Y. (1968) The red-backed vole on Rishiri island. *Hoppo Ringyo* (Northern Forestry) **20**: 169-172. (in Japanese)
- Misawa, E. (1979) Change in the food habits of the red fox, *Vulpes vulpes schrencki* Kishida according to habitat conditions. *Journal of the Mammalogical Society of Japan* **7**:311-320. (in Japanese with English summary).
- Modi, W.S. and R. Gamperl (1989) Chromosomal banding comparisons among American and European red-backed mice, genus *Clethrionomys*. *Zeitschrift f r Säugertierkunde* **54**: 141-152.
- Musser, G. G. and M. D. Carleton. (1993) Family Muridae. pp. 501-755. In Wilson, Don E. and D.-A. M. Reeder (eds.) *Mammal Species of the World*, 2nd edn. Smithsonian Institution Press, Washington and London.
- Nakata, K. (1984) Factors affecting litter size in the red-backed vole, *Clethrionomys rufocanus bedfordiae*, with special emphasis on population phase. *Researches on Population Ecology* **26**: 221-234.
- Nakata, K. (1986) *Handbook for vole census methods and control*. Hokkaido Shinrin-Hozen Kyokai, Sapporo. (in Japanese)
- Nakata, K. (1987) Effects of spring-removal on the populations of small rodents. *Japanese Journal of Applied Entomology and Zoology* **31**: 97-91 (in Japanese with English summary)
- Nakata, K. (1989) Regulation of reproductive rate in a cyclic population of the red-backed vole, *Clethrionomys rufocanus bedfordiae*. *Researches on Population Ecology* **31**:185-209.
- Nakata, K. (1995) Microhabitat selection in two sympatric species of voles *Clethrionomys rex* and *Clethrionomys rufocanus bedfordiae*. *Journal of the Mammalogical Society of Japan* **20**:135-142.
- Nakata, K. (1998) *Historical review of forecasting program for vole density*. Forest Protection Supplement, Hokkaido Shinrin-Hozen Kyokai, Sapporo. (in Japanese)
- Niethammer, J. and F. Krapp (eds. 1982) *Handbuch der Säugetiere Europas. Vol.1 & 2*. Akademische Verlags, Wiesbaden.
- Nowak, R. M. and J. L. Paradiso (1983) *Walker's mammals of the world. Vol. 2*. 4th edn. The Johns Hopkins University Press, Baltimore.
- Ognev, S. I. (1950) *Mammals of the USSR and Adjacent Countries. Vol. 7 Rodents*. Israel Prog. Sci. Trans. (English version published in 1964)
- Obara, Y., H. Kusakabe, K. Miyakoshi and S. Kawada (1995) Revised karyotypes of the Japanese northern red-backed vole, *Clethrionomys rutilus mikado*. *Journal of the Mammalogical Society of Japan* **20**: 125-133.
- Ota, K. (1956) The Muridae of the islands adjacent to Hokkaido. *Memoirs of the Faculty of Agriculture, Hokkaido University* **2**: 123-136. (in Japanese with English abstract)
- Ota, K. (1958a) Studies on the interspecific relationship of murid rodents. III. Habitat segregation a small tree stand. *Japanese Journal of Ecology* **8**: 149-156. (in Japanese)
- Ota, K. (1958b) The relationship between forest damage by voles and climate in Hokkaido. *Nonezumi* (Voles and Mice) **29**: 1-2. (in Japanese)
- Ota, K. (1968) Studies on the ecological distribution of the murid rodents in Hokkaido. *Research Bulletins of the College Experiment Forests, College of Agriculture, Hokkaido University* (Sapporo) **26**: 223-295. (in Japanese with English summary)
- Ota, K.(ed.) (1984) *Study on wild murid rodents in Hokkaido*. Hokkaido University Press, Sapporo. (in Japanese)
- Ota, K. and J. Fujikura (1959) Report on the ecology of voles in the pilot forests (No. 6). *Juhyo* **9**(7): 12-25. (in Japanese)
- Ota, K. and E. W. Jameson, Jr. (1961) Ecological relationships and economic importance of Japanese Microtinae. *Ecology* **42**: 184-186.
- Ota, K., S. Takatsu and H. Abe (1959) Fluctuations of numbers in small mammals at Mt. Moiwa, Sapporo. I. Seasonal fluctuations of small mammal populations. *Memoirs of the Faculty of Agriculture, Hokkaido University* (Sapporo) **3**: 49-69. (in Japanese with English summary)
- Rauschert, K. (1963) Sexuelle Affinität zwischcen Arten und Unterarten von Rattelnäusen (*Clethrionomy*). *Biologisches Zentralblatt* **82**: 653-664.
- Saitoh, T. (1981) Control of female maturation in high density populations of the red-backed vole, *Clethrionomys rufocanus bedfordiae*. *Journal of Animal Ecology* **50**:79-87.

- Saitoh, T. (1987) A time series and geographical analysis of population dynamics of the red-backed vole in Hokkaido, Japan. *Oecologia* **73**:382-388.
- Saitoh, T. (1989) Effects of added food on some attributes of an enclosed vole population. *Journal of Mammalogy* **70**:772-782.
- Saitoh, T. (1990) Lifetime reproductive success in reproductively suppressed female voles. *Researches on Population Ecology* **32**:391-406.
- Saitoh, T. (1991) The effects and limits of territoriality on population regulation in grey red-backed voles, *Clethrionomys rufocanus bedfordiae*. *Researches on Population Ecology* **33**:367-386.
- Saitoh, T. and A. Nakatsu (1997) Impact of forest plantation on the community of small mammals in Hokkaido, Japan. *Mammal Study* **22**: 27-38.
- Saitoh, T. and K. Takahashi (1998) The role of vole populations in prevalence of the parasite (*Echinococcus multilocularis*) in foxes. *Researches on Population Ecology* **40**: 97-105.
- Saitoh, T., N. C. Stenseth, and O. N. Bjørnstad. 1997. Density dependence in fluctuating grey-sided vole populations. *Journal of Animal Ecology* **66**:14-24.
- Saitoh, T., N. C. Stenseth, and O. N. Bjørnstad (1998a) Density-dependence in voles and mice: a comparative study. *Ecology* (in press)
- Saitoh, T., N. C. Stenseth, and O. N. Bjørnstad (1998b) The population dynamics of the vole, *Clethrionomys rufocanus*, in Hokkaido. *Researches on Population Ecology* **40**: 61-76.
- Semenov-Tjan-Shanskij, O. E. (1970) Ciklicnost' v populacijah lesnyh polevok. *Bjull Mv. o.-va isp. periody, otd. biol.* **75**: 11-26. (in Russian)
- Skar, H.-J., A. Hagen and E. Jystbye (1971) The bank vole *Clethrionomys glareolus* (Schreber, 1780) in south Norwegian mountain area. *Norwegian Journal of Zoology* **19**: 261-266.
- Stenseth, N. C. (ed.) (1985a) *Clethrionomys* biology: population dynamics, dispersal, reproduction and social structure. *Annales Zoologici Fennici* **22**: 205-395.
- Stenseth, N. C. (1985b) Geographical distribution of *Clethrionomys* species. *Annales Zoologici Fennici* **22**: 215-219.
- Stenseth, N. C., O. N. Bjørnstad and T. Saitoh (1996) A gradient from stable to cyclic populations of *Clethrionomys rufocanus* in Hokkaido, Japan. *Proceedings of the Royal Society of London B* **263**: 1117-1126.
- Thomas, O. (1905a) Abstract. *Proceedings of the Zoological Society of London* **23**: 18-19.
- Thomas, O. (1905b) The Duke of Bedford's zoological expedition in eastern Asia. I. List of mammals obtained by Mr. M. P. Anderson in Japan. *Proceedings of the Zoological Society of London* 1905: 331-363.
- Tokuda, M. (1935) *Neoschizomys*, a new genus of Microtinae from Shikotan, a South Kuril Islands. *Memoirs College Science Kyoto Imperial University, Series B* **10**: 241-250.
- Tokuda (1941) A revised monograph of the Japanese and Manchou-Korean Muridae. *Transaction of the Biogeographical Society of Japan* **4**: 1-155.
- Ueda, M. (1961) A preliminary report on outbreaks of vole populations in the Daisetsu District Office. *Transaction of the Meeting in Hokkaido Branch of the Japanese Forestry Society* **10**: 134-136. (in Japanese)
- Ueda, M., S. Higuchi, B. Igarashi, M. Maeda, T. Kuwahata, K. Ota, H. Abe, Y. Fujimaki, J. Fujikura and T. Takayasu (1966) Historical review of studies on the Bedford's red-backed vole, *Clethrionomys rufocanus bedfordiae* (Thomas). *Bulletin of the Government Forest Experiment Station (Tokyo)* **191**: 1-100. (in Japanese)
- Uraguchi, K., T. Saitoh, N. Kondo and H. Abe (1987) Food habits of the feral mink (*Mustela vison* Schreber) in Hokkaido. *Journal of the Mammalogical Society of Japan* **12**:57-67.
- Viitala, J. (1971) Age determination in *Clethrionomys rufocanus* (Sundevall). *Annales Zoologici Fennici* **8**: 63-67.
- Viitala, J. (1977) Social organization in cyclic subarctic populations of the voles *Clethrionomys rufocanus* (Sund.) and *Microtus atrestis* (L.). *Annales Zoologici Fennici* **14**:53-93.
- Vorontsov, N. N., E. A. Lyapunova, Yu. M. Borissov and V. E. Dovgal (1980) Variability of sex chromosomes in mammals. *Genetica* **52/53**: 361-372.
- Wada, N. (1993) Dwarf bamboos affect the regeneration of zoochorous trees by providing habitats to acorn-feeding rodents. *Oecologia* **94**:403-407.
- Wakana, S., M. Sakaizumi, K. Tsuchiya, M. Asakawa, S. H. Han, K. Nakata and H. Suzuki (1996) Phylogenetic implications of variation in rDNA and mtDNA in red-backed voles collected in Hokkaido, Japan and Korea. *Mammal Study* **21**: 12-25.
- Whitney, P. (1977) Seasonal maintenance and net production of two sympatric species of subarctic microtine rodents. *Ecology* **28**: 314-325.
- Wilson, Don E. and D.-A. M. Reeder (eds.) (1993) *Mammal species of the world*, 2nd edn. Smithsonian Institute Press, Washington and London.
- Yoneda, M. (1979) Prey preference of the red fox,

- Vulpes vulpes schrencki* Kishida (Carnivora: Canidae), on small rodents. *Applied Entomology and Zoology* **14**:28-35.
- Ylönen, H. and J. Viitala (1986) Social organization and habitat use of introduced populations of the vole *Clethrionomys rufocanus* (Sund) in central Finland. *Zeitschrift für Säugetierkunde* **52**: 354-363.
- Yoccoz, N G., K. Nakata, N. C. Stenseth and T. Saitoh (1998) The demography of the vole *Clethrionomys rufocanus*: from mathematical and statistical models to further field studies. *Researches on Population Ecology* **40**: 107-121.
- Yoneda, M. (1982) Influence of red fox predation on a local population of small rodents I. Population fluctuation in small rodents. *Applied Entomology and Zoology* **17**:8-19.
- Yoneda, M. (1983) Influence of red fox predation on a local population of small rodents. II. Seasonal changes in predation pressure, prey preference and predation effect. *Applied Entomology and Zoology* **18**:1-10.
- Yoneda, M. and K. Nakata (1984) Natural enemies. pp. 159-185. In Ota, K.(ed.) *Study on wild murid rodents in Hokkaido*. Hokkaido University Press, Sapporo. (in Japanese)
- Yoneda, M., H. Abe and H. Nakao (1979) Winter food habits of the Yezo Ural Owl *Strix uralensis japonica* in a wind shelter-belt. *Miscellaneous Reports of Yamashina Institute of Ornithology* **11**:49-53. (in Japanese with English summary)
- Yoshida, I., Y. Obara and N. Matsuoka (1989) Phylogenetic relationship among seven taxa of the Japanese microtine voles revealed by karyological and biochemical techniques. *Zoological Science* **6**: 409-420.
- Zimmermann, V. K. (1964) Zur Säugetier-Fauna China. *Mitteilungen aus dem Zoologischen Museum in Berlin* **40**: 87-140. (in German)
- Zimmerman, V. K. (1965) Art-Hybriden bei *Rattus*. *Zeitschrift für Säugetierkunde* **30**: 315-320. (in German)

Appendix. A list of scientific names of genera, species and subspecies in *Clethrionomys rufocanus* according to historical order described.

-
- ***1811** *Hypudaeus* Illiger [Type: *Mus lemmus* Linnaeus, 1758; *Mus amphibius* Linnaeus 1758; and *Mus arvalis* Pallas, 1779] *Hypudaeus* is a synonym of *Lemmus* Link, 1795
- ***1846** *Hypudaeus rufocanus* Sundevall [Type locality: Lappmark, Sweden]
- 1850** *Clethrionomys* Tilesius [Type: *Mus rutilus* Pallas, 1779]
- ***1874** *Evotomys* Coues [Type: *Mus rutilus* Pallas, 1779]
- ***1881** *Arvicola wosnessenskii* Polyakov [Type locality: Kamchatka, Russia]
- ***1881** *Arvicola rufocanus* var. *sibirica* Polyakov [Type locality: Kamchatka, Russia]
- 1884** *Arvicola rufocanus* var. *kamtschaticus* Lataste [Type locality: Kamchatka, Russia] *nom. nud.*
- 1900** *Craseomys* Miller [Type: *Hypudaeus rufocanus* Sundevall, 1846]
- 1903** *Evotomys* (*Craseomys*) *latastei* Allen [Type locality: Gichiga, Kamchatka, Russia] Allen replaced *Arvicola rufocanus* var. *kamtschaticus* Lataste by this form, because *Arvicola kamtschaticus* (*nec.*) Polyakov was preoccupied as a primary homonym.
- 1905** *Evotomys bedfordiae* Thomas [Type locality: Shinshinotsu, near Sapporo, Hokkaido, Japan]
- 1917** *Evotomys* (*Craseomys*) *irkutensis* Ognev [Type locality: Irkutsk Province, Siberia, Russia]
- 1922** *Evotomys kolymensis* Ognev [Type locality: Berezova near Sredne Kolymsk, Russia]
- 1924** *Craseomys rufocanus bargusiniensis* Turov [Type locality: Bargusin Reserve, northeastern shore, Lake Baikal, Russia]
- 1928** *Evotomys* (*Craseomys*) *arsenjevi* Dukelski [Type locality: Vladivostok, Ussuri Bay, Russia]
- 1930** *Clethrionomys rufocanus kurilensis* Kishida [Type locality: not described] *nom. nud.*
- 1932** *Clethrionomys rufocanus kurilensis* Tokuda [Type locality: Paramushir Island, Kuril Islands, Japan]
- 1935** *Neoaschizomys sikotanensis* Tokuda [Type locality: Shikotan Island, the southern part of Kuril Islands, Japan]
- 1949** *Neoaschizomys sikotanensis akkessi* Imaizumi [Type locality: Daikoku Islet, off Hokkaido, Japan] *Neoaschizomys*
- ***1984** *Clethrionomys rufocanus bromleyi* Kostenko [Type locality: Sakhalin, Russia] *nom. nud.*
- 1993** *Clethrionomys rufocanus changbaishanensis* Jang, Ma and Luo [Type locality: Anty county, Jinlin Province, China]
-

* not seen the original papers.